

ABSTRACTS

List of Talks

Vladimir Bogachev: Nonlinear Kantorovich problems and Hausdorff distances between plans	2
Alexander Bufetov: The Gaussian multiplicative chaos for the sine-process	2
Marat Burnashev: On Stein’s lemma in a general case	3
Alexander Gushchin: Old and new results on sets of joint distributions of certain pairs of stochastic processes	5
Rustam Ibragimov: New approaches to robust inference on market (non-)efficiency, volatility clustering and nonlinear dependence	5
Alexey Khartov: Quasi-infinitely divisible distributions	6
Vladimir Lotov: Exact formulas in some boundary crossing problems for integer-valued random walks	7
Alexey Naumov: On Dirichlet boundary crossing	7
Alexander Nazarov: Spectral analysis and L_2 -small ball asymptotics of some FBM-like processes	8
Ernst Presman: An Inventory Model with Markov Chain Modulated Commodity Prices and long-run average costs	8
Evgeny Prokopenko: Multi-Normex approach for evaluating the sum of heavy tailed random vectors	9
Alexander Sakhanenko: On first-passage times for generalized random walks	11
Stanislav Shaposhnikov: The superposition principle for Fokker–Planck–Kolmogorov equations	12
Irina Shevtsova: Comparison of central and noncentral Lyapounov fractions with application to Berry–Esseen bounds for Poisson random sums	12
Natalia Stepanova: Sparse signal recovery with Subbotin noise	13
Alexander Tikhomirov: Limit theorems for spectrum of random graphs	13
Vladimir Ulyanov: Inference via randomized test statistics	14
Anatoly Vershik: Uniqueness of invariant Markov numerations and tilings of posets	14
Elena Yarovaya: Martingale method for studying branching random walks	15

Vladimir Bogachev: Nonlinear Kantorovich problems and Hausdorff distances between plans

Moscow State University and Higher School of Economics
vibogach@mail.ru

We discuss recent investigations of nonlinear Kantorovich problems of optimal transportation involving cost functions that can depend on transportation plans, in particular, on conditional measures of transportation plans. Some existence results will be presented. In addition, we consider Hausdorff distances on sets of transportation plans associated with the Kantorovich and Kantorovich–Rubinshtein metrics on spaces of probability measures. Such Kantorovich-type Hausdorff distances admit simple bounds via distances between marginal distributions. Some open problems in this area are also mentioned. More details on this direction can be found in the forthcoming paper [1].

References

- [1] V.I. Bogachev. The Kantorovich problem of optimal transportation of measures: new directions of research. *Russian Math. Surveys.*, 77 (2022) (in press).

Alexander Bufetov: The Gaussian multiplicative chaos for the sine-process

Steklov Mathematical Institute and Aix-Marseille University, France
bufetov@mi.ras.ru

To almost every realization of the sine-process one naturally assigns a random entire function, the analogue of the Euler product for the sine, the scaling limit of ratios of characteristic polynomials of a random matrix. The main result of the talk is that the square of the absolute value of our random entire function converges to the Gaussian multiplicative chaos. As a corollary, one obtains that almost every realization with one particle removed is a complete and minimal set for the Paley-Wiener space, whereas if two particles are removed, then the resulting set is a zero set for the Paley-Wiener space. Quasi-invariance of the sine-process under compactly supported diffeomorphisms of the line plays a key rôle.

In joint work with Qiu, the Patterson-Sullivan construction is used to interpolate Bergman functions from a realization of the determinantal point process with the Bergman kernel, in other words, by the Peres-Virág theorem, the zero set of a random series with independent complex Gaussian entries. The invariance of the zero set under the isometries of the Lobachevsky plane plays a key rôle. Conditional measures of the determinantal point process with the Bergman kernel are found explicitly (cf. arXiv:2112.15557, Dec. 2021).

Marat Burnashev: On Stein's lemma in a general case

Institute for Information Transmission Problems, RAS, Moscow
burn@iitp.ru

Introduction. Let \mathbf{P} and \mathbf{Q} be probability measures on a measurable space $(\mathcal{X}, \mathcal{B})$. We consider testing of simple hypothesis \mathcal{H}_0 versus simple alternative \mathcal{H}_1 , based on observation $\mathbf{x} \in \mathcal{X}$:

$$\mathcal{H}_0 : \mathbf{x} \sim \mathbf{P}, \quad \mathcal{H}_1 : \mathbf{x} \sim \mathbf{Q}. \quad (1)$$

If for testing a decision (acceptance) region $\mathcal{D} \in \mathcal{X}$ is chosen, such that

$$\mathbf{x} \in \mathcal{D} \Rightarrow \mathcal{H}_0, \quad \mathbf{x} \notin \mathcal{D} \Rightarrow \mathcal{H}_1, \quad (2)$$

then type-I error probability $\alpha(\mathcal{D})$ and type-II error probability $\beta(\mathcal{D})$ are defined by formulas, respectively,

$$\alpha(\mathcal{D}) = \mathbf{P}(\mathbf{x} \notin \mathcal{D} | \mathcal{H}_0), \quad \beta(\mathcal{D}) = \mathbf{P}(\mathbf{x} \in \mathcal{D} | \mathcal{H}_1). \quad (3)$$

We consider the case when type-I error probability α , $0 < \alpha < 1$, is fixed, and we are interested in the minimal possible type-II error probability $\beta(\alpha)$

$$\beta(\alpha) = \inf_{\mathcal{D}: \alpha(\mathcal{D}) \leq \alpha} \beta(\mathcal{D}). \quad (4)$$

The corresponding optimal decision region $\mathcal{D}(\alpha)$ to this problem is given by Neyman-Pearson lemma. We will need the following notion.

Definition 1. For probability distributions \mathbf{P} and \mathbf{Q} on $(\mathcal{X}, \mathcal{B})$ the function $D(\mathbf{P}||\mathbf{Q})$ (Kullback–Leibler distance or divergence) is defined as

$$D(\mathbf{P}||\mathbf{Q}) = \mathbf{E}_{\mathbf{P}} \ln \frac{d\mathbf{P}}{d\mathbf{Q}}(\mathbf{x}) \geq 0, \quad (5)$$

where the expectation is taken over the measure \mathbf{P} .

In order to describe Stein's lemma and its relation to the value $\beta(\alpha)$ via the function $D(\mathbf{P}||\mathbf{Q})$, we consider first the case $\mathcal{X} = \mathbf{R}^n$, when \mathbf{x} from (1) has the form $\mathbf{x} = \mathbf{x}_n = (x_1, \dots, x_n) \in \mathbf{R}^n$. We assume that the sample \mathbf{x}_n consists of i.i.d. random variables x_i , $i = 1, \dots, n$ with a distribution \mathbf{P} (in the case of \mathcal{H}_0), or with a distribution \mathbf{Q} (in the case of \mathcal{H}_1):

$$\mathcal{H}_0 : x_i \sim \mathbf{P}, \quad \mathcal{H}_1 : x_i \sim \mathbf{Q}, \quad i = 1, \dots, n. \quad (6)$$

If in model (6) $\mathbf{P} \neq \mathbf{Q}$, then $D(\mathbf{P}||\mathbf{Q}) > 0$ and the value $\beta(\alpha)$ decreases exponentially as $n \rightarrow \infty$. Moreover, we have [3, Theorem 3.3]

$$\lim_{\alpha \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} \ln \beta(\alpha) = -D(\mathbf{P}||\mathbf{Q}). \quad (7)$$

Relation (7) is called Stein's lemma [1, 2].¹ In the case of i.i.d. random variables, its proof can be found in [3, Theorem 3.3], [4, Theorem 12.8.1]. It is natural to expect that relation (7) holds not only for i.i.d. random variables as in model (6), but in much more general cases. Some particular

¹The reference [1] is due to remarks from [2, p. 18] and [3, Theorem 3.3] with attribute to the unpublished paper of C. Stein.

analog of formula (7) have already appeared for cases of stationary Gaussian [5] and Poisson [6] random processes (with time T instead of n).

Relation (7) is oriented to the case when type-I and type-II errors imply very different losses, and we wish to minimize type-II error probability $\beta(\alpha)$. The case is quite popular in many applications (see, e.g., [5] and references therein). In this paper, for general model (1) some non-asymptotic generalization and strengthening of relation (7) is derived.

Main result. In order to simplify formulation of results and to avoid pathological cases, we assume that the following assumption is satisfied:

I. Probability measures \mathbf{P} and \mathbf{Q} on $(\mathcal{X}, \mathcal{B})$ are equivalent and Radon-Nikodim derivative $d\mathbf{P}/d\mathbf{Q}(\mathbf{x})$ is defined, positive and finite for $\mathbf{x} \in \mathcal{X}$. The main result is the following

Theorem. *If the assumption I holds, then the minimal possible $\beta(\alpha)$, $0 < \alpha < 1$, satisfies the bounds*

$$\begin{aligned} \ln \beta(\alpha) &\geq -\frac{D(\mathbf{P}||\mathbf{Q}) + h(\alpha)}{1 - \alpha}, \\ h(\alpha) &= -\alpha \ln \alpha - (1 - \alpha) \ln(1 - \alpha), \end{aligned} \quad (8)$$

and

$$\ln \beta(\alpha) \leq -D(\mathbf{P}||\mathbf{Q}) + \mu_0(\alpha), \quad (9)$$

where $\mu_0(\alpha)$ is the minimal value μ_0 , satisfying the relation

$$\mathbf{P}_{\mathbf{P}} \left\{ \ln \frac{d\mathbf{P}}{d\mathbf{Q}}(\mathbf{x}) \leq D(\mathbf{P}||\mathbf{Q}) - \mu_0 \right\} \leq \alpha. \quad (10)$$

With some loss of accuracy we can simplify upper bound (9), using an upper bound for the value $\mu_0(\alpha)$. For example, using Chebyshev inequality, we get

Corollary. *The minimal possible $\beta(\alpha)$, satisfies also the upper bound*

$$\ln \beta(\alpha) \leq -D(\mathbf{P}||\mathbf{Q}) + \alpha^{-1/2} r_1(\mathbf{P}, \mathbf{Q}), \quad (11)$$

where

$$r_1(\mathbf{P}, \mathbf{Q}) = \left\{ \mathbf{E}_{\mathbf{P}} \left[\ln \frac{d\mathbf{P}}{d\mathbf{Q}}(\mathbf{x}) \right]^2 - D^2(\mathbf{P}||\mathbf{Q}) \right\}^{1/2}. \quad (12)$$

Note that all bounds (8), (9) and (11) are pure analytical relations without any limiting operations.

References

- [1] C. Stein. Information and comparison of experiments (*unpublished*).
- [2] H. Chernoff. Large-sample theory: parametrics case. *Ann. Math. Statist.*, 27:1-22, 1956.
- [3] S. Kullback. *Information Theory and Statistics*. Wiley, New York, 1958.
- [4] T. M. Cover and J. A. Thomas. *Elements of Information Theory*. Wiley, New York, 1991.
- [5] W. Zhang and H. V. Poor. On minimax robust detection of stationary Gaussian signals in white Gaussian noise. *IEEE Trans. Inf. Theory*, 57,6:3915-3924, 2011.
- [6] M. V. Burnashev. On Neyman-Pearson minimax detection of Poisson process intensity. *Statistical Inference for Stochastic Processes*. 24,1:211-221, 2021.

Alexander Gushchin: Old and new results on sets of joint distributions of certain pairs of stochastic processes

Steklov Mathematical Institute
gushchin@mi-ras.ru

Based on the joint work with Mikhail Nedoshivin (Lomonosov Moscow State University)

The aim of this talk is twofold. First, we recall known results describing the sets of joint distributions of certain pairs of stochastic processes, namely, a martingale and its maximum [5], and an increasing process and its compensator [2]. We also show how these results are connected with the Skorokhod embedding problem [?]. Second, we show how to adapt the Perkins construction [4], [1] in the Skorokhod embedding problem to the above problems and demonstrate some new results.

References

- [1] A. M. G. Cox and D. G. Hobson. An optimal Skorokhod embedding for diffusions. *Stochastic Process. Appl.*, 111(1):17-39, 2004.
- [2] A. A. Gushchin. The joint law of terminal values of a nonnegative submartingale and its compensator. *Theory Probab. Appl.*, 62(2):216–235, 2018.
- [3] A. A. Gushchin. The joint law of a max-continuous local submartingale and its maximum. *Theory Probab. Appl.*, 65(4):545–557, 2021.
- [4] E. Perkins. The Cereteli–Davis solution to the H^1 -embedding problem and an optimal embedding in Brownian motion. In: *Seminar on stochastic processes, 1985*, Progress in Probability and Statistics 12, Birkhäuser, Boston, 1986, 172–223.
- [5] L. C. G. Rogers. The joint law of the maximum and terminal value of a martingale. *Probab. Theory Relat. Fields*, 95(4):451–466, 1993.

Rustam Ibragimov: New approaches to robust inference on market (non-)efficiency, volatility clustering and nonlinear dependence

Imperial College Business School, UK
ribragim@gmail.com

Rasmus S. Pedersen (Department of Economics, Inniversitu of Copenhagen) and Anton Skrobotov (RANEPA)

Many financial and economic variables, including financial returns, exhibit nonlinear dependence, heterogeneity and heavy-tailedness. These properties may make problematic the analysis of (non-) efficiency and volatility clustering in economic and financial markets using traditional approaches that appeal to asymptotic normality of sample autocorrelation functions of returns and their squares. This paper presents new approaches to deal with the above problems. We provide the results that motivate the use of measures of market (non-)efficiency and volatility clustering based on (small)

powers of absolute returns and their signed versions. We further provide new approaches to robust inference on the measures in the case of general time series, including GARCH-type processes. The approaches are based on robust t-statistics tests based on the results by Bakirov and Szekely (Zap. Nauch. Sem. POMI, 2005; Journal of Mathematical Sciences, 2006) and new results on their applicability are presented. In the approaches, parameter estimates (e.g., estimates of measures of nonlinear dependence) are computed for groups of data, and the inference is based on t-statistics in the resulting group estimates. This results in valid robust inference under heterogeneity and dependence assumptions satisfied in real-world financial markets. Numerical results and empirical applications confirm the advantages and wide applicability of the proposed approaches.

Alexey Khartov: Quasi-infinitely divisible distributions

Smolensk State University
alexeykharov@gmail.com

The new class of *quasi-infinitely divisible probability laws* is a natural and rather wide extension of the class of infinitely divisible laws. According to the definition, a probabilistic law on the real line is called a quasi-infinitely divisible if its characteristic function admits the Lévy-Khinchin representation with a real shift parameter and with some non-monotonic (in general case) spectral function having bounded variation on the whole real line. Examples of such laws can be found in the well-known classical monographs by Gnedenko, Kolmogorov [1], and by Linnik, Ostrovsky [2]. However, the definition and the corresponding class were introduced only in 2011 by Lindner and Sato within some problems for stochastic processes. Further, these laws find some applications in the number theory, finance and physics. The first detailed analysis of the class of quasi-infinitely divisible laws was recently made by Lindner, Pan and Sato in [3]. So now this class is actually studied. In the talk, we review basic facts concerning this class and we also present new results for it, which were obtained in [4, 5, 6].

References

- [1] B.V. Gnedenko and A.N. Kolmogorov. *Limit Distributions for Sums of Independent Random Variables*. Addison-Wesley, Cambridge, 1954.
- [2] Yu.V. Linnik and I.V. Ostrovskii. *Decomposition of Random Variables and Vectors*. Transl. Math. Monog., Vol. 48, AMS, Providence, Rhode Island, 1977.
- [3] A. Lindner, L. Pan and K. Sato. On quasi-infinitely divisible distributions. *Trans. Amer. Math. Soc.*, 370:8483–8520, 2018.
- [4] I.A. Alexeev and A.A. Khartov. Spectral representations of characteristic functions of discrete probability laws. *arXiv:2101.06038 (Accepted in Bernoulli)*, 2022.
- [5] A.A. Khartov. A criterion of quasi-infinite divisibility for discrete laws. *Stat.&Prob. Letters*, 185:109436, 2022.
- [6] A.A. Khartov. On weak convergence of quasi-infinitely divisible laws. *arXiv:2204.13667*, 2022.

Vladimir Lotov: Exact formulas in some boundary crossing problems for integer-valued random walks

Sobolev Institute of Mathematics and Novosibirsk State University
lotov49@mail.ru

Let X_1, X_2, \dots be i.i.d. integer-valued random variables,

$$S_n = X_1 + \dots + X_n, \quad n \geq 1, \quad S_0 = 0.$$

We suppose that the distribution of X_1^+ belongs to the class \mathbf{K} that contains distributions of bounded random variables, convolutions of geometric distributions and their mixtures.

For arbitrary $b \geq 1$, introduce

$$\tau = \tau(b) = \inf\{n \geq 1 : S_n \geq b\}, \quad \chi = \chi(b) = S_\tau - b.$$

Suppose that $\tau = \infty$ if $S_n < b$ for all n . In this case, the value of χ remains undefined.

The main goal is to find exact formulas:

- (i) for $P(\chi(b) = k)$, $k \geq 1$;
- (ii) for the renewal function $E\tau(b)$ if $EX_1 > 0$;
- (iii) for $P(\sup S_n \geq b)$ if $EX_1 < 0$.

We start with the factorization representation for the double moment generating function $E(z^\tau \mu^\chi; \tau < \infty)$, analyze its structure and demonstrate the way to inverse it.

This work was supported by the Russian Science Foundation (grant no. 22-21-00396).

Alexey Naumov: On Dirichlet boundary crossing

Higher School of Economics
anaumov@hse.ru

Based on the joint work with D. Belomestny (Duisburg-Essen University), P. Menard (Otto von Guericke University), E. Moulines (Ecole Polytechnique), S. Samsonov (HSE University), Y. Tang (Deepmind), D. Tiapkin (HSE University) and M. Valko (Deepmind)

In the talk we will provide upper and lower bounds on the deviation of a linear function of Dirichlet random vector (Dirichlet boundary crossing). The second result is devoted to a tight Gaussian lower bound for the distribution of a linear function of Dirichlet random vector. In particular, we extend the normal approximation-based lower bound for Beta distributions by [1]. These problems surprisingly arise in the estimation of regret for many modern reinforcement learning algorithms.

References

- [1] D. Alfers and H. Dinges. A normal approximation for beta and gamma tail probabilities. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete.*, 65:399–420, 1984.

Alexander Nazarov: Spectral analysis and L_2 -small ball asymptotics of some FBM-like processes

PDMI RAS and St. Petersburg State University
al.il.nazarov@gmail.com

We study the spectral problems for some integro-differential equations arising in the theory of Gaussian processes similar to the fractional Brownian motion. We generalize the method of Chigansky–Kleptsyna [1] and obtain the two-term eigenvalues asymptotics for such equations. Application to the small ball probabilities in L_2 -norm is given.

The talk is based on the papers [2], [3] and on some recent results in preparation. The research is supported by the RFBR grant 20-51-12004.

References

- [1] P. Chigansky and M. Kleptsyna. Exact Asymptotics in Eigenproblems for Fractional Brownian Covariance Operators. *Stochastic Processes and Their Applications*. 2018. Vol. 128, No. 6, pp. 2007–2059.
- [2] A.I. Nazarov. Spectral asymptotics for a class of integro-differential equations arising in the theory of fractional Gaussian processes. *Communications in Contemporary Mathematics*. 2020. DOI 10.1142/S0219199720500492, pp. 1–25.
- [3] A.I. Nazarov. Some lemmata on the perturbation of the spectrum. *Russian Journal of Mathematical Physics*. 2020. Vol. 27, No. 3, pp. 378–381.

Ernst Presman: An Inventory Model with Markov Chain Modulated Commodity Prices and long-run average costs

MSE MGU, CEMI RAS
presman@cemi.rsi.ru

The following inventory model was considered in [1]: A manufacturer (firm) uses for “production” a “commodity” (resource), which is consumed with unit intensity. The price of the commodity follows a stochastic process, modulated by a continuous time Markov chain with a finite number of states and known transition rates. The firm can buy this commodity at the current price or use “stored” one. The storage cost is proportional to the storage level. The goal of the firm is to minimize the total discounted performance costs. The existence of an optimal “threshold” strategy was proven. The threshold strategy is defined by a vector of levels specifying the minimal commodity amount to keep for a given price. An algorithm was presented to find an optimal “thresholds” and the corresponding value functions. It was shown that as a rule such strategy is unique and all situations were described when it is not unique. In typical optimization problems in continuous time involving Bellman (optimality) equation, a smooth pasting of the first derivatives

of the value functions is used. A special feature of the considered model was that: the smooth pasting takes place for the second derivative of the value functions, it is convenient to conduct reasoning not with the price of the game itself, but with its derivative, and the Bellman equation itself is convenient to write out for the imbedded chain.

In this paper we consider the same model but with long-run average costs. To investigate this case, we aim the discount coefficient to zero. As a result, we obtain an analogue of the canonical triplet, well-known in discrete-time Markov chain control problems, which consists of optimal control, the optimal value of the long-run average cost functional and the invariant functional. We prove the existence of optimal threshold strategy and show that the algorithm for finding the optimal thresholds and values of the derivative of the invariant functional does not differ from the case of discounting, but to find the invariant functional itself, it is necessary to analyze the features when the stock level tends to infinity. As a rule, the optimal stationary strategy is again unique, but there may again be cases of non-uniqueness, while in the pre-limit case the optimal strategy is unique.

This work was supported by Russian Science Foundation (Project 20-68-47030)

References

- [1] E.L. Presman, I.M. Sonin. Model upravlinia zapasami s tsenami na syrio, zavisyashchimi ot tsepi Markova s neprerivnim vremenem. *Zhournal Novoi Ekonomicheskoi Asociaii*, presented at 2022.

Evgeny Prokopenko: Multi-Normex approach for evaluating the sum of heavy tailed random vectors

Novosibirsk State University and Sobolev Institute of Mathematics
e.prokopenko@ngs.ru

Based on the joint work with Marie Kratz (ESSEC Business School, CREAR, France)

Looking for the most accurate possible evaluation of the distribution of the sum of random variables, or vectors, or processes, with unknown distributions, has always been a classical problem in the probabilistic and statistical literature, with various answers depending on the given framework and on the specific application in view. On one hand, (uni- or multivariate) Central Limit Theorems (CLT) or Functional ones prove, under finite variance for the sum components or/and additional conditions, the asymptotic Gaussian behavior of the sum with some rate of convergence, focusing on the 'body' of the distribution. When considering heavy-tailed marginal distributions, Generalized CLT with the convergence to stable distributions, handle the case of infinite variance (see e.g. [1], [2], and references therein), while, in the case of finite variance, an alternative way is to consider trimmed sums, removing extremes from the sample, to improve the rate of convergence; see e.g. [3], [4] and references therein.

When interested in tail distributions, CLTs may give poor results, especially when considering heavy tails. That is why different approaches have been developed, among which: (i) large deviation theorems; (ii) extreme value theorems (EVT) focusing on the tail only; (iii) hybrid distributions combining (asymptotic) distributions for both the main and extreme behaviors when considering

independent random variables (see e.g. [5, 6, 7, 8]; we use the name given in [7] for this type of hybrid distribution/method/approach, namely *Normex* distribution/method/approach.

Recall briefly the idea of Normex (for 'Norm(al)-Ex(tremes)') method. It consists of rewriting the sum of random variables as the sum of their ordered statistics, and splitting it into two main parts, a trimmed sum removing the extremes, and the extremes. Using that the trimmed sum of the first $n - k - 1$ ordered statistics is conditionnally independent of the k largest order statistics, given the $(n - k)$ -th order statistics, we can express the distribution of the sum, integrating w.r.t. to the $(n - k)$ -th order statistics and using a CLT for the conditional trimmed sum, and an EVT one for the k largest order statistics. Note that a benefit of Normex approach is that it does not require any condition on the existence of moments, as the CLT applies on truncated random variables.

It is natural to extend the normex approach to a multivariate framework. With this goal of proposing a multi-normex method and distribution, we consider iid random vectors $\mathbf{X}_1, \dots, \mathbf{X}_n$, with parent random vector \mathbf{X} having a heavy-tailed d -dimensional distribution $F_{\mathbf{X}}$ and density $f_{\mathbf{X}}$ (when existing). Note that there are different ways to define multivariate extremes. The chosen way in this paper is w.r.t. the norm $\|\cdot\|$ in \mathbb{R}^d , meaning that the ordered (w.r.t. the norm) vector of $(\mathbf{X}_1, \dots, \mathbf{X}_n)$, denoted by $(\mathbf{X}_{(1)}, \dots, \mathbf{X}_{(n)})$, satisfies

$$\|\mathbf{X}_{(1)}\| \leq \|\mathbf{X}_{(2)}\| \leq \dots \leq \|\mathbf{X}_{(n)}\|.$$

Aiming at proving the benefit of using a multi-normex distribution for a better fit of the whole (unknown) heavy-tailed distribution $F_{\mathbf{X}}$, assuming $F_{\mathbf{X}}$ heavy-tailed in the sense of $\|\mathbf{X}\| \in \mathcal{RV}_{-\alpha}$, we focus analytically on the case $\alpha \in (2; 3]$ (when $\|\mathbf{X}\|$ has a finite second moment, but no third moment) to compare the rates of convergence when using the CLT and the multi-normex approach, respectively. Note our focus on heavy tailed distributions (i.e. distributions belonging to the max domain of attraction of Fréchet), where the impact of using Normex distribution will be much stronger than in the light tail case (because of the one big jump principle), in particular for risk analysis and management. We prove that the normex approach leads, as expected, to a better speed of convergence for evaluating the distribution of the sum than the CLT does, for such type of heavy-tailed distributions. When varying the fatness of the tail measured by $\alpha > 0$, we draw this comparison numerically, using geometrical multivariate quantiles.

References

- [1] Samorodnitsky, G. and M. S. Taqqu. *Stable non-Gaussian Random Processes: Stochastic Models with Infinite Variance*. Chapman and Hall, 1994.
- [2] Petrov, V. V. *Sums of Independent Random Variables*. Ergebnisse der Mathematik und ihrer Grenzgebiete, Springer, 1975.
- [3] Mori, T. On the limit distributions of lightly trimmed sums. *Mathematical Proceedings of the Cambridge Philosophical Society*, 96.3:507–516, 1984.
- [4] Hahn, M. G. Sums. *Trimmed Sums and Extremes*. Progress in Probability. Ed. by M. G. Hahn, D. M. Mason, and D. C. Weiner. Vol. 23. Birkhäuser Boston, 1991.
- [5] Csörgö, S., E. Haeusler, and D. M. Mason. A probabilistic approach to the asymptotic distribution of sums of independent, identically distributed random variables. *Advances in Applied Mathematics*, 9.3:259–333, 1988.

- [6] Zaliapin, I. V., Y. Y. Kagan, and F. P. Schoenberg. Approximating the Distribution of Pareto Sums. *Pure and Applied Geophysics*, 162:1187–1228, 2005.
- [7] Kratz, M. Normex, a new method for evaluating the distribution of aggregated heavy tailed risks. Application to risk measures. *Extremes*, 17.4:661–691, 2014.
- [8] Müller, U. K. Refining the central limit theorem approximation via extreme value theory. *Statistics and Probability Letters*, 155:1–7, 2019.

Alexander Sakhanenko: On first-passage times for generalized random walks

Sobolev Institute of Mathematics, Novosibirsk, Russia
 aisakh@mail.ru

The talk is based on the joint works with D. Denisov (University of Manchester, UK) and V. Wachtel (University of Bielefeld, Germany). The research was funded by RFBR and DFG according to the research project 20-51-12007.

Let $S(t)$ be a random process defined for all $t \geq 0$ with $S(0) = 0$ and let $g(t)$ be a non-random function on $[0, \infty)$. Introduce the random variable

$$\tau := \inf \{t > 0 : S(t) \leq g(t)\} = \inf \{t > 0 : S(t) - g(t) \leq 0\},$$

equal to the first moment of the top-down crossing of the level $g(t)$ by our process $S(t)$. We consider in the talk the asymptotic behavior of the upper tail $\mathbf{P}(\tau > T)$ as $T \rightarrow \infty$.

For several classes of processes we are going to obtain conditions under which we have asymptotical formulas of the following type:

$$\mathbf{P}(\tau > T) \sim \frac{U(T)}{\sqrt{T}} \quad \text{as } T \rightarrow \infty, \tag{1}$$

for some slowly varying functions $U(T)$.

Two such classes of processes may be found in papers [1] and [2]. In these cases

$$U(T) := \sqrt{\frac{2}{\pi}} \mathbf{E}[S(\alpha_T) - g(\alpha_T); \tau > \alpha_T] \tag{2}$$

for specially chosen stopping times α_T such that $\alpha_T/T \rightarrow 0$ in probability as $T \rightarrow \infty$.

References

- [1] D. Denisov, A. Sakhanenko, V. Wachtel. First-passage times for random walks with non-identically distributed increments. *Ann. Probab.*, 46:3313–3350, 2018.
- [2] A. Shelepova, A. Sakhanenko. On the asymptotics of the probability to stay above a non-increasing boundary for a non-homogeneous compound renewal process. *Sib. Electronic Math. Reports*, 18:1667–1688, 2021.

Stanislav Shaposhnikov: The superposition principle for Fokker–Planck–Kolmogorov equations

Moscow State University
starticle@mail.ru

We discuss the superposition principle for probability solutions to the Cauchy problem for the Fokker–Planck–Kolmogorov equation, according to which such a solution μ_t with initial distribution ν is represented by a probability measure P_ν on the path space such that P_ν solves the corresponding martingale problem and μ_t is the one-dimensional distribution of P_ν at time t . We present new sufficient conditions for the superposition principle to be fulfilled in the case of unbounded coefficients. Moreover the superposition principle for equations on a domain is considered.

References

- [1] V.I. Bogachev, N.V. Krylov, M. Röckner, S.V. Shaposhnikov, *Fokker–Planck–Kolmogorov Equations*. Amer. Math. Soc., Rhode Island, Providence, 2015.
- [2] V.I. Bogachev, M. Röckner, S.V. Shaposhnikov. On the Ambrosio–Figalli–Trevisan superposition principle for probability solutions to Fokker–Planck–Kolmogorov equations, *J. Dynamics Differ. Equ.*, 33, 715–739, 2021.

Irina Shevtsova: Comparison of central and noncentral Lyapounov fractions with application to Berry–Esseen bounds for Poisson random sums

Moscow State University
ishevtsova@cs.msu.ru

Based on the joint work with Vladimir Makarenko (Moscow State University)

For a nondegenerate r.v. X with finite $E|X|^3$ let us denote

$$L_1(X) = \frac{E|X|^3}{(EX^2)^{3/2}}, \quad L_0(X) = L_1(X - EX) = \frac{E|X - EX|^3}{(DX)^{3/2}}.$$

We find an explicit expression for the least upper bound

$$H(t) := \sup_{EX=t\sqrt{EX^2}} \frac{L_1(X)}{L_0(X)}$$

for every $t \in (-1, 1)$, as well as its unconditional supremum

$$\sup_{-1 < t < 1} H(t) = \frac{1}{4} \sqrt{17 + 7\sqrt{7}} = 1.48997\dots$$

The numerical result $\sup_{-1 < t < 1} H(t) < 1.49$ was previously obtained in [2] with the help of a computer.

Furthermore, we apply these results to the Berry–Esseen-type inequality for Poisson random sums [1, 3]

$$\Delta_\lambda := \sup_{x \in \mathbf{R}} \left| \mathbb{P} \left(\frac{S_\lambda - \mathbb{E}S_\lambda}{\sqrt{\mathbb{D}S_\lambda}} < x \right) - \Phi(x) \right| \leq 0.3031 \frac{L_1(X)}{\sqrt{\lambda}}, \quad \lambda > 0,$$

where $S_\lambda = X_1 + \dots + X_{N_\lambda}$, N_λ being a Poisson random variable with expectation λ , independent of i.i.d. r.v.'s $\{X_k\}_{k \geq 1}$, to deduce its analogue involving the non-central Lyapounov fraction

$$\Delta_\lambda \leq 0.3031 \cdot H(t) \frac{L_0(X)}{\sqrt{\lambda}} \leq 0.4517 \frac{L_0(X)}{\sqrt{\lambda}}, \quad \lambda > 0,$$

for every distribution of X_1 with fixed $\mathbb{E}X_1/\sqrt{\mathbb{E}X_1^2} = t \in (-1, 1)$. A similar upper bound, uniform with respect to t , was previously obtained in [2].

References

- [1] V. Korolev and I. Shevtsova. An improvement of the Berry–Esseen inequality with applications to Poisson and mixed Poisson random sums. *Scand. Actuar. J.*, 2012:81–105, 2012.
- [2] V. Yu. Korolev, I. G. Shevtsova and S. Ya. Shorgin. On the Berry–Esseen type inequalities for Poisson random sums. *Inform. Primen.*, 5:64–66, 2011.
- [3] I. G. Shevtsova. On the absolute constants in the Berry–Esseen-type inequalities. *Dokl. Math.*, 89:378–381, 2014.

Natalia Stepanova: Sparse signal recovery with Subbotin noise

Carleton University, Canada
 nstep@math.carleton.ca

Based on the joint work with Joshua Miller (Carleton University)

We study the problem of high-dimensional variable selection in a sparse sequence model with Subbotin noise. We derive the regions of exact and almost full variable selection with respect to a Hamming loss function, and propose adaptive procedures that achieve both types of selection. The proposed procedures are shown to be optimal in the asymptotically minimax sense. Our results augment previous work in this area.

Alexander Tikhomirov: Limit theorems for spectrum of random graphs

Komi Science Center of Ural Branch of RAS
 sasha-tikh@yandex.ru

We consider the limit of empirical spectral distribution of Laplace matrix and adjacency matrix of generalized random graphs. Applying Stieltjes transform method we proof under general conditions that limit spectral distribution of adjacency matrix is Wigner law and empirical spectral distribution of Laplace matrices convergence to the free convolution of Wigner law and normal law.

Vladimir Ulyanov: Inference via randomized test statistics

Moscow State University and Higher School of Economics
vulyan@gmail.com

Based on the joint work with Nikita Puchkin (HSE University and IITP RAS)

We show that external randomization may enforce the convergence of test statistics to their limiting distributions in particular cases. This results in a sharper inference. Our approach is based on a central limit theorem for weighted sums. We apply our method to a family of rank-based test statistics and a family of phi-divergence test statistics (see e.g.[1]) and prove that, with high probability with respect to the external randomization, the randomized statistics converge at the rate $O(1/n)$ (up to some logarithmic factors) to the limiting chi-square distribution in Kolmogorov metric. See the details in [2].

References

- [1] Y. Fujikoshi and V.V. Ulyanov. *Non-Asymptotic Analysis of Approximations for Multivariate Statistics*. SpringerBriefs in Statistics. Springer Nature, Singapore, 2020.
- [2] N. Puchkin and V. Ulyanov. Inference via Randomized Test Statistics. *ArXiv*, 2112.06583, 2021.

Anatoly Vershik: Uniqueness of invariant Markov numerations and tilings of posets

PDMI RAS and St. Petersburg State University
avershik@gmail.com

Partially based on the joint work with Fedya Petrov

We introduce a new class of probabilistic-combinatorial problems: to describe a Markov chain which monotonically numerates the elements of a countable poset, or, for a continual poset, monotonically tiles it by segments. We look for the “most random” way, i.e., the way with the maximal entropy. Such problems have very different applications, from Young diagrams statistics to the spectra of random matrices, etc. In most cases the answer is unique up to intensity of numeration or tiling, and the obtained Markov chains are related to Bernoulli schemes in a simple but non-trivial manner. Even the simplest examples which we discuss give interesting and unusual answers, and the explicit computation of transition probabilities is quite difficult.

Elena Yarovaya: Martingale method for studying branching random walks

Moscow State University
yarovaya@mech.math.msu.su

Based on the joint work with Natalia Smorodina (St. Petersburg Department of Steklov Mathematical Institute of Russian Academy of Sciences)

We consider a continuous-time branching random walk on a multidimensional lattice \mathbb{Z}^d , $d \in \mathbb{N}$, in which particles can die and produce offspring at any point on the lattice, see, for example, [1]. Let the transport of particles on \mathbb{Z}^d be given by a symmetric, homogeneous and irreducible random walk. A branching intensity at a point $x \in \mathbb{Z}^d$ tends to zero as $\|x\| \rightarrow \infty$. Moreover, an additional condition is satisfied for the parameters of the branching random walk, which guarantees exponential growth in time of the average number of particles at each point $x \in \mathbb{Z}^d$. Under these assumptions, we prove the limit theorem about the mean square convergence of the normalized number of particles at an arbitrary fixed point $x \in \mathbb{Z}^d$, as $t \rightarrow \infty$. The proof is based on the approximation of the normalized number of particles by a nonnegative martingale [2].

The research was supported in part by the Russian Foundation for Basic Research (RFBR), Project No. 20-01-00487.

References

- [1] E. Yarovaya. *Branching Random Walks in non Homogeneous Environments*. Center of Applied Investigations of the Faculty of Mechanics and Mathematics of the Moscow State University, Moscow, Russian, 2007.
- [2] N. Smorodina and E. Yarovaya. A Martingale Method for Studying Branching Random Walks. *Russian Math. Surveys*, 5, 2022 (in print).