

INDIAN STATISTICAL INSTITUTE
ST. PETERSBURG DEPARTMENT OF STEKLOV MATHEMATICAL INSTITUTE OF THE RUSSIAN ACADEMY OF
SCIENCES

LABORATORY ON ALGEBRAIC TRANSFORMATION GROUPS AT HSE UNIVERSITY

Indian-Russian Conference
Polynomial Rings and Group Actions

Saint Petersburg, Russia
May 27 – June 1, 2024

ABSTRACTS

Contents

<i>Ivan Arzhantsev.</i> Tits-type alternative for \mathbb{G}_a -generated groups of automorphisms . . .	3
<i>Roman Avdeev.</i> Root subgroups on horospherical varieties	4
<i>Prosenjit Das.</i> Translates of a line	5
<i>Nikhilesh Dasgupta.</i> On commuting locally nilpotent derivations and isotropy subgroups in $k[X, Y, Z]$	6
<i>Neena Gupta.</i> On the generalisation of linear planes of Sathaye and Russel	7
<i>Animesh Lahiri.</i> On affine forms	8
<i>Swapnil A. Lokhande.</i> A survey on finite generation of kernel of locally nilpotent derivations of polynomial rings	9
<i>Ananya Pal.</i> On rigidity of Pham-Brieskorn surfaces	10
<i>Alexander Perepechko.</i> Structure of connected nested automorphism groups	11
<i>Victor Petrov.</i> Tits construction and the Rost invariant	12
<i>Anton Shafarevich.</i> On modified Derksen and Makar-Limanov invariants	13
<i>Anastasia Stavrova.</i> Local triviality of principal G -bundles over the relative projective line	14
<i>Yulia Zaitseva.</i> Algebraic monoid structures on the affine 3-space	15

Tits-type alternative for \mathbb{G}_a -generated groups of automorphisms

Ivan Arzhantsev

HSE University

arjantsev@hse.ru

In 1972, Tits has proved that any subgroup of the general linear group over a field of characteristic zero either contains a nonabelian free subgroup or is a finite extension of a solvable group. In the first part of the talk we survey known facts on the Tits alternative for groups of regular and birational automorphisms of algebraic varieties.

The second part of the talk is based on joint works with Mikhail Zaidenberg. Let X be an affine algebraic variety defined over an algebraically closed field of characteristic zero and \mathbb{G}_a be the additive group of the ground field. Consider a subgroup H of the automorphism group $\text{Aut}(X)$ generated by a finite collection of \mathbb{G}_a -subgroups. We conjecture that either H contains a nonabelian free subgroup or H is a unipotent affine algebraic group. We prove this conjecture when X is an affine toric variety and the \mathbb{G}_a -subgroups are normalized by the acting torus. Also we show that if X is a complex affine surface then either H contains a nonabelian free subgroup or H is a metabelian unipotent algebraic group.

The research is supported by the RSF-DST grant 22-41-02019.

Root subgroups on horospherical varieties

Roman Avdeev
HSE University
ravdeev@hse.ru

In the study of automorphism groups of toric varieties, a key role is played by one-parameter additive groups normalized by the acting torus. Such subgroups are called root subgroups and each of them is uniquely determined by its weight, called a Demazure root of the corresponding toric variety. Moreover, the set of all Demazure roots admits an explicit combinatorial description in terms of the fan defining the toric variety, and this description is especially simple in the case where the variety is affine.

In the setting of arbitrary connected reductive groups acting on algebraic varieties, a natural generalization of toric varieties is given by spherical varieties. A spherical variety is an algebraic variety X equipped with an action of a connected reductive group G in such a way that a Borel subgroup B of G has a dense open orbit in X . A proper generalization of root subgroups for spherical varieties is given by one-parameter additive groups normalized by B , which are called B -root subgroups.

The most accessible for study class of spherical varieties is given by horospherical ones. In the talk we shall discuss B -root subgroups on horospherical varieties.

The talk is based on joint works of the speaker with I. Arzhantsev and with V. Zhgoon. The work is supported by the RSF-DST grant 22-41-02019.

Translates of a line
Prosenjit Das
Department of Mathematics,
Indian Institute of Space Science and Technology
prosenjit.das@iist.ac.in

It is readily apparent that any coordinate of $k^{[2]}$ is a line of $k^{[2]}$ and any translate of a coordinate of $k^{[2]}$ is again a coordinate of $k^{[2]}$. In a landmark result, Abhyankar-Moh and Suzuki, established that over a field k of characteristic zero any line of $k^{[2]}$ is a coordinate of $k^{[2]}$, known as the Epimorphism Theorem. Consequently, it follows that in a polynomial algebra in two indeterminates over a field of characteristic zero, lines and coordinates are synonymous; and translate of a line is a line too.

However, it is a well-established fact that over fields of positive characteristic, the equivalence between lines and coordinates no longer holds true; a line need not be a coordinate. This leads to an intriguing, yet unresolved problem posed by Sathaye which asks whether any translate of line remains a line.

In this talk, which forms part of a collaborative work with Animesh Lahiri, shall discuss on the captivating question posed by Sathaye.

On commuting locally nilpotent derivations and isotropy subgroups in $k[X, Y, Z]$
Nikhilesh Dasgupta
Department of Mathematics,
GITAM University Bengaluru
`its.nikhilesh@gmail.com`

For a locally nilpotent derivation on an affine algebraic variety, its isotropy subgroup is defined to be the group of algebraic automorphisms which commute with the locally nilpotent derivation. Let k be a field of characteristic zero. In this talk, I would like to discuss the properties of commuting locally nilpotent derivations on the polynomial ring $k[X, Y, Z]$ and their isotropy subgroups.

This talk is based on recent works with S. Gayfullin and A. Lahiri.

On the generalisation of linear planes of Sathaye and Russell
Neena Gupta
Theoretical Statistics and Mathematics Unit Indian Statistical Institute
rnanina@gmail.com

In this talk we shall present several examples where the Abhyankar-Sathaye Conjecture for linear hyperplanes has been confirmed.

On affine forms
Animesh Lahiri
GITAM University Bengaluru
alahiri@gitam.edu

Let k be a field with algebraic closure \bar{k} and A be a k -algebra. We say that A is an \mathbb{A}^n -form over k if $A \otimes_k \bar{k} = \bar{k}^{[n]}$. It is well-known that separable \mathbb{A}^1 -forms are trivial (i.e., $k^{[1]}$) and that there exist non-trivial purely inseparable \mathbb{A}^1 -forms over fields of positive characteristic. An extensive study of such algebras was made by T. Asanuma. T. Kambayashi established that separable \mathbb{A}^2 -forms over a field k are trivial. Over any field of positive characteristic, the non-trivial purely inseparable \mathbb{A}^1 -forms can be used to give examples of non-trivial \mathbb{A}^n -forms for any integer $n > 1$. However, the problem of existence of non-trivial separable \mathbb{A}^3 -forms over a field is still open in general.

Now let R be a ring containing a field k . An R -algebra A is said to be an \mathbb{A}^n -form over R with respect to k if $A \otimes_k \bar{k} = (R \otimes_k \bar{k})^{[n]}$, where \bar{k} denotes the algebraic closure of k . A.K. Dutta has investigated separable \mathbb{A}^1 -forms over any ring R containing a field k and as a corollary obtained that the separable \mathbb{A}^2 -forms over a PID containing \mathbb{Q} are the trivial ones.

In this talk, we will discuss a partial result on separable \mathbb{A}^3 -forms over a field k and show an extension of the results on \mathbb{A}^2 -forms to any one-dimensional Noetherian \mathbb{Q} -algebra and to any \mathbb{Q} -algebra having a fixed point free locally nilpotent derivation.

**A survey on finite generation of kernel of locally nilpotent derivations of
polynomial rings**

Swapnil A. Lokhande

Department of Humanities and Sciences,

IIIT Vadodara

`swapnil@iiitvadodara.ac.in`

Hilbert's fourteenth problem, in context of locally nilpotent derivations, asks whether kernel of a locally nilpotent derivation (LND) of a finitely generated k -algebra is finitely generated. In this talk, we give a survey of all the known results and discuss results on LNDs of $R[X, Y, Z]$, where R is a UFD and finitely generated k -algebra.

On rigidity of Pham-Brieskorn surfaces
Ananya Pal
Stat-Math Unit, Indian Statistical Institute
palananya1995@gmail.com

An irreducible affine variety \mathbb{V} over a field k is called rigid if it admits no non-trivial \mathbb{G} -action. Any \mathbb{G} -action on \mathbb{V} has an equivalent formulation as an exponential map on the corresponding coordinate ring $k[\mathbb{V}]$. Thus a k -algebra A is said to be rigid if it has no non-trivial exponential map. We define a k -algebra A to be stably rigid if for any $n \geq 0$ and for any exponential map ϕ on the polynomial ring $A[X_1, \dots, X_n]$, we have $A \subseteq A[X_1, \dots, X_n]^\phi$.

For any three integers $a, b, c \geq 1$, the rigidity of Pham-Brieskorn surfaces,

$$B_{(a,b,c)} := \frac{k[X, Y, Z]}{(X^a + Y^b + Z^c)},$$

has been studied by several experts including G. Freudenburg, A. J. Crachiola, S. Maubach, when k is an algebraically closed field of characteristic zero.

In this talk we are going to discuss the rigidity of Pham-Brieskorn rings over an arbitrary field k of characteristic $p \geq 0$. We give some sufficient conditions on (a, b, c) for which any Pham-Brieskorn domain $B_{(a,b,c)}$ is rigid. This gives an alternative approach to show that there does not exist any non-trivial exponential map on $k[X, Y, Z, T]/(X^m Y + T^{p^r q} + Z^{p^e}) = k[x, y, z, t]$, for $m, q > 1$, $p \nmid mq$ and $e > r \geq 1$, which fixes y , a crucial result used in [1] to show that the Zariski Cancellation Problem does not hold for the affine 3-space.

We also provide a sufficient condition for $B_{(a,b,c)}$ to be stably rigid.

This is a joint work with Neena Gupta.

References

- [1] Neena Gupta, *On the cancellation problem for the affine space \mathbb{A}^3 in characteristic p* , Invent. Math. **195** (2014) 279–288.

Structure of connected nested automorphism groups

Alexander Perepechko

IITP RAS/HSE University

aperepechko@hse.ru

A nested group is an increasing union of algebraic groups. It is well known that any algebraic subgroup of the automorphism group $\text{Aut}(X)$ of an affine variety X is closed with respect to the ind-topology. The closedness of connected nested subgroups in $\text{Aut}(X)$ is an open question.

In this talk, we answer this question positively. Moreover, we establish the structure of nested subgroups of $\text{Aut}(X)$. In the course of the proof, we describe maximal nested unipotent subgroups of $\text{Aut}(X)$ and their relation to \mathbb{G}_a^k -actions on X .

The work is supported by the RSF-DST grant 22-41-02019.

Tits construction and the Rost invariant

Victor Petrov

SPbSU/PDMI RAS

victorapetrov@googlemail.com

Simple Lie algebras over an algebraically closed field of characteristic 0 are described by Dynkin diagrams. Over a non-closed field, the same Dynkin diagram can correspond to many simple algebras, so it is interesting to study constructions of simple Lie algebras and invariants that make it possible to recognize their isomorphism or reflect some of their properties. One such construction of exceptional (i.e., types E_6 , E_7 , E_8 , F_4 or G_2) Lie algebras was proposed by Jacques Tits; the Jordan algebra and an alternative algebra are given as input, and the output is a Lie algebra, and all real forms of Lie algebras can be constructed in this way. One of the most useful invariants (with meaning in the third Galois cohomology group) was constructed by Markus Rost. We show that a Lie algebra of (outer) type E_6 is obtained by the Tits construction if and only if the Rost invariant is a pure symbol. As an application of this result we prove a Springer-type theorem for an E_6 -homogeneous manifold.

On modified Derksen and Makar-Limanov invariants

Anton Shafarevich

MSU University/HSE University

shafarevich.a@gmail.ru

The talk is based on joint work with Sergey Gaifullin.

The modified Derksen invariant $\text{HD}^*(X)$ of an affine algebraic variety X is the subalgebra in $\mathbb{K}[X]$ generated by kernels of all locally nilpotent derivations of $\mathbb{K}[X]$ with slices. If there is a locally nilpotent derivation of $\mathbb{K}[X]$ with a slice then $X \simeq Y \times \mathbb{A}^1$ where Y is an affine variety. In my talk we will show that there are three possibilities: A) $\text{HD}^*(X) = \mathbb{K}[X]$; B) $\text{HD}^*(X)$ is a proper infinitely generated subalgebra; C) $\text{HD}^*(X) = \mathbb{K}[Y]$. We give examples for each case, and also provide sufficient conditions for the variety Y so that the variety $X = Y \times \mathbb{A}^1$ belongs to one of the types.

The work is supported by the RSF-DST grant 22-41-02019.

Local triviality of principal G -bundles over the relative projective line

Anastasia Stavrova

PDMI RAS

`anastasia.stavrova@gmail.com`

Let U be a semilocal noetherian integral scheme. Let G be a reductive group scheme over U . We show that a principal G -bundle over the relative projective line \mathbb{P}_U^1 is Zariski locally trivial, once it is trivial at the infinity section. The talk is based on the joint work with Ivan Panin.

Algebraic monoid structures on the affine 3-space

Yulia Zaitseva
HSE University
yzaytseva@hse.ru

The talk is based on joint work with Ivan Arzhantsev and Roman Avdeev. We have completed the classification of algebraic monoids on the affine 3-space. The most challenging case was that of non-commutative monoids with a group of invertible elements of rank one, which we reduced to the corresponding commutative case.

The work is supported by the RSF-DST grant 22-41-02019.