

# Convolution Polynomials

Donald E. Knuth  
Computer Science Department  
Stanford, California 94305-2140

**Abstract.** The polynomials that arise as coefficients when a power series is raised to the power  $x$  include many important special cases, which have surprising properties that are not widely known. This paper explains how to recognize and use such properties, and it closes with a general result about approximating such polynomials asymptotically.

A family of polynomials  $F_0(x), F_1(x), F_2(x), \dots$  forms a *convolution family* if  $F_n(x)$  has degree  $\leq n$  and if the convolution condition

$$F_n(x+y) = F_n(x)F_0(y) + F_{n-1}(x)F_1(y) + \dots + F_1(x)F_{n-1}(y) + F_0(x)F_n(y)$$

holds for all  $x$  and  $y$  and for all  $n \geq 0$ . Many such families are known, and they appear frequently in applications. For example, we can let  $F_n(x) = x^n/n!$ ; the condition

$$\frac{(x+y)^n}{n!} = \sum_{k=0}^n \frac{x^k}{k!} \frac{y^{n-k}}{(n-k)!}$$

is equivalent to the binomial theorem for integer exponents. Or we can let  $F_n(x)$  be the binomial coefficient  $\binom{x}{n}$ ; the corresponding identity

$$\binom{x+y}{n} = \sum_{k=0}^n \binom{x}{k} \binom{y}{n-k}$$

is commonly called *Vandermonde's convolution*.

How special is the convolution condition? *Mathematica* will readily find all sequences of polynomials that work for, say,  $0 \leq n \leq 4$ :

```
F[n_,x_] := Sum[f[n,j] x^j, {j,0,n}]/n!
conv[n_] := LogicalExpand[Series[F[n,x+y], {x,0,n}, {y,0,n}]
  == Series[Sum[F[k,x] F[n-k,y], {k,0,n}], {x,0,n}, {y,0,n}]]
Solve[Table[conv[n], {n,0,4}],
  [Flatten[Table[f[i,j], {i,0,4}, {j,0,4}]]]]
```

*Mathematica* replies that the  $F$ 's are either identically zero or the coefficients of  $F_n(x) = (f_{n0} + f_{n1}x + f_{n2}x^2 + \dots + f_{nn}x^n)/n!$  satisfy

$$\begin{aligned} f_{00} &= 1, & f_{10} &= f_{20} = f_{30} = f_{40} = 0, \\ f_{22} &= f_{11}^2, & f_{32} &= 3f_{11}f_{21}, & f_{33} &= f_{11}^3, \\ f_{42} &= 4f_{11}f_{31} + 3f_{21}^2, & f_{43} &= 6f_{11}^2f_{21}, & f_{44} &= f_{11}^4. \end{aligned}$$

This allows us to choose  $f_{11}$ ,  $f_{21}$ ,  $f_{31}$ , and  $f_{41}$  freely.

Suppose we weaken the requirements by asking only that the convolution condition hold when  $x = y$ . The definition of `conv` then becomes simply

```
conv[n_] := LogicalExpand[Series[F[n, 2x], {x, 0, n}]
  == Series[Sum[F[k, x] F[n-k, x], {k, 0, n}], {x, 0, n}]]
```

and we discover that the same solutions occur. In other words, the weaker requirements imply that the strong requirements are fulfilled as well.

In fact, it is not difficult to discover a simple rule that characterizes all “convolution families.”  
Let

$$F(z) = 1 + F_1 z + F_2 z^2 + F_3 z^3 + \cdots$$

be any power series with  $F(0) = 1$ . Then the polynomials

$$F_n(x) = [z^n] F(z)^x$$

form a convolution family. Conversely, every convolution family arises in this way or is identically zero. (Here the notation  $[z^n] \text{expr}$  stands for what *Mathematica* calls `Coefficient[expr, z, n]`.)

*Proof.* Let  $f(z) = \ln F(z) = f_1 z + f_2 z^2/2! + f_3 z^3/3! + \cdots$ . It is easy to verify that the coefficient of  $z^n$  in  $F(z)^x$  is indeed a polynomial in  $x$  of degree  $\leq n$ , because  $F(z)^x = e^{x f(z)} = \exp(x f_1 z + x f_2 z^2/2! + x f_3 z^3/3! + \cdots)$  expands to the power series

$$\sum_{k_1, k_2, k_3, \dots \geq 0} x^{k_1 + k_2 + k_3 + \cdots} \frac{f_1^{k_1} f_2^{k_2} f_3^{k_3} \cdots}{1!^{k_1} k_1! 2!^{k_2} k_2! 3!^{k_3} k_3! \cdots} z^{k_1 + 2k_2 + 3k_3 + \cdots};$$

when  $k_1 + 2k_2 + 3k_3 + \cdots = n$  the coefficient of  $z^n$  is a polynomial in  $x$  with terms of degree  $k_1 + k_2 + k_3 + \cdots \leq n$ . This construction produces a convolution family because of the rule for forming coefficients of the product  $F(z)^{x+y} = F(z)^x F(z)^y$ .

Conversely, suppose the polynomials  $F_n(x)$  form a convolution family. The condition  $F_0(0) = F_0(0)^2$  can hold only if  $F_0(x) = 0$  or  $F_0(x) = 1$ . In the former case it is easy to prove by induction that  $F_n(x) = 0$  for all  $n$ . Otherwise, the condition  $F_n(0) = 2F_n(0)$  for  $n > 0$  implies that  $F_n(0) = 0$  for  $n > 0$ . If we equate coefficients of  $x^k$  on both sides of

$$F_n(2x) = F_n(x)F_0(x) + F_{n-1}(x)F_1(x) + \cdots + F_1(x)F_{n-1}(x) + F_0(x)F_n(x),$$

we now find that the coefficient  $f_{nk}$  of  $x^k$  in  $n! F_n(x)$  is forced to have certain values based on the coefficients of  $F_1(x), \dots, F_{n-1}(x)$ , when  $k > 1$ , because  $2^k f_{nk}$  occurs on the left and  $2f_{nk}$  on the right. The coefficient  $f_{n1}$  can, however, be chosen freely. Any such choice must make  $F_n(x) = [z^n] \exp(x f_{11} z + x f_{21} z^2/2! + x f_{31} z^3/3! + \cdots)$ , by induction on  $n$ .

**Examples.** The first example mentioned above,  $F_n(x) = x^n/n!$ , comes from the power series  $F(z) = e^z$ ; the second example,  $F_n(x) = \binom{x}{n}$ , comes from  $F(z) = 1 + z$ . Several other power series

are also known to have simple coefficients when we raise them to the power  $x$ . If  $F(z) = 1/(1-z)$ , for instance, we find

$$[z^n](1-z)^{-x} = \binom{-x}{n}(-1)^n = \binom{x+n-1}{n}.$$

It is convenient to use the notations

$$\begin{aligned} x^{\underline{n}} &= x(x-1) \dots (x-n+1) = x!/(x-n)! \\ x^{\overline{n}} &= x(x+1) \dots (x+n-1) = \Gamma(x+n)/\Gamma(x) \end{aligned}$$

for falling factorial powers and rising factorial powers. Since  $\binom{x}{n} = x^{\underline{n}}/n!$  and  $\binom{x+n-1}{n} = x^{\overline{n}}/n!$ , our last two examples have shown that the polynomials  $x^{\underline{n}}/n!$  and  $x^{\overline{n}}/n!$  form convolution families, corresponding to  $F(z) = 1+z$  and  $F(z) = 1/(1-z)$ . Similarly, the polynomials

$$F_n(x) = \frac{x(x-s)(x-2s) \dots (x-(n-1)s}{n!}$$

form a convolution family corresponding to  $(1+sz)^{1/s}$  when  $s \neq 0$ .

The cases  $F(z) = 1+z$  and  $F(z) = 1/(1-z)$  are in fact simply the cases  $t=0$  and  $t=1$  of a general formula for the binomial power series  $\mathcal{B}_t(z)$ , which satisfies

$$\mathcal{B}_t(z) = 1 + z \mathcal{B}_t(z)^t.$$

When  $t$  is any real or complex number, exponentiation of this series is known to yield

$$[z^n] \mathcal{B}_t(z)^x = \binom{x+tn}{n} \frac{x}{x+tn} = \frac{x(x+tn-1) \dots (x+tn-n+1)}{n!};$$

see, for example, [Graham et al 1989, section 7.5, example 5], where a combinatorial proof is given.

The special cases  $t=2$  and  $t=-1$ ,

$$\begin{aligned} \mathcal{B}_2(z) &= \frac{1 - \sqrt{1-4z}}{2z} = 1 + z + 2z^2 + 5z^3 + 14z^4 + 42z^5 + \dots, \\ \mathcal{B}_{-1}(z) &= \frac{1 + \sqrt{1+4z}}{2} = 1 + z - z^2 + 2z^3 - 5z^4 + 14z^5 - \dots, \end{aligned}$$

in which the coefficients are the Catalan numbers, arise in numerous applications. For example,  $\mathcal{B}_2(z)$  is the generating function for binary trees, and  $\mathcal{B}_1(-z)$  is the reciprocal of  $\mathcal{B}_2(z)$ . We can get identities in trigonometry by noting that  $\mathcal{B}_2((\frac{1}{2} \sin \theta)^2) = \sec^2(\theta/2)$ . Furthermore, if  $p$  and  $q$  are probabilities with  $p+q=1$ , it turns out that  $\mathcal{B}_2(pq) = 1/\max(p, q)$ . The case  $t=1/2$ ,

$$\mathcal{B}_{1/2}(z) = \left( \frac{z + \sqrt{4+z^2}}{2} \right)^2 = 1 + z + \frac{z^2}{2} + \frac{z^3}{2^3} - \frac{z^5}{2^7} + \frac{2z^7}{2^{11}} - \frac{5z^9}{2^{15}} + \frac{14z^{11}}{2^{19}} - \dots,$$

is another interesting series in which the Catalan numbers can be seen. The convolution polynomials in this case are the ‘‘central factorials’’  $x(x + \frac{n}{2} - 1)^{\underline{n-1}}/n!$  [Riordan 1968, section 6.5], also called Steffensen polynomials [Roman and Rota 1978, example 6].

The convolution formula corresponding to  $\mathcal{B}_t(z)$ ,

$$\binom{x+y+tn}{n} \frac{x+y}{x+y+tn} = \sum_{k=0}^n \binom{x+tk}{k} \frac{x}{x+tk} \binom{y+t(n-k)}{n-k} \frac{y}{y+t(n-k)}$$

is a rather startling generalization of Vandemonde's convolution; it is an identity for all  $x, y, t$ , and  $n$ .

The limit of  $\mathcal{B}_t(z/t)^t$  as  $t \rightarrow \infty$  is another important function  $T(z)/z$ ; here

$$T(z) = \sum_{n \geq 1} \frac{n^{n-1}}{n!} z^n = z + z^2 + \frac{3z^3}{2} + \frac{8z^4}{3} + \frac{125z^5}{24} + \dots$$

is called the *tree function* because  $n^{n-1}$  is the number of labeled, rooted trees. The tree function satisfies

$$T(z) = ze^{T(z)},$$

and we have the corresponding convolution family

$$[z^n] \left( \frac{T(z)}{z} \right)^x = [z^n] e^{xT(z)} = \frac{x(x+n)^{n-1}}{n!}.$$

The related power series

$$1 + zT'(z) = \frac{1}{1-T(z)} = \sum_{n \geq 0} \frac{n^n z^n}{n!} = 1 + z + 2z^2 + \frac{9z^3}{2} + \frac{32z^4}{3} + \frac{625z^5}{24} + \dots$$

defines yet another convolution family of importance: We have

$$[z^n] \frac{1}{(1-T(z))^x} = \frac{t_n(x)}{n!},$$

where  $t_n(x)$  is called the *tree polynomial* of order  $n$  [Knuth and Pittel 1989]. The coefficients of  $t_n(x) = t_{n1}x + t_{n2}x^2 + \dots + t_{nn}x^n$  are integers with combinatorial significance; namely,  $t_{nk}$  is the number of mappings of an  $n$ -element set into itself having exactly  $k$  cycles.

A similar but simpler sequence arises from the coefficients of powers of  $e^{ze^z}$ :

$$n! [z^n] e^{xze^z} = \sum_{k=0}^n \binom{n}{k} k^{n-k} x^k.$$

The coefficient of  $x^k$  is the number of *idempotent* mappings of an  $n$ -element set into itself, having exactly  $k$  cycles [Harris and Schoenfeld 1967].

If the reader still isn't convinced that convolution families are worthy of detailed study, well, there's not much hope, although another example or two might clinch the argument. Consider the power series

$$e^{e^z-1} = \sum \frac{b_n z^n}{n!} = 1 + \frac{z}{1!} + \frac{2z^2}{2!} + \frac{5z^3}{3!} + \frac{15z^4}{4!} + \frac{52z^5}{5!} + \dots;$$

these coefficients  $b_n$  are the so-called *Bell numbers*, the number of ways to partition sets of size  $n$  into subsets. For example, the five partitions that make  $b_3 = 5$  are

$$\{1, 2, 3\}, \quad \{1\}\{2, 3\}, \quad \{1, 2\}\{3\}, \quad \{1, 3\}\{2\}, \quad \{1\}\{2\}\{3\}.$$

The corresponding convolution family is

$$[z^n] e^{(e^z - 1)x} = \frac{\{0\}^n + \{1\}^n x + \{2\}^n x^2 + \cdots + \{n\}^n x^n}{n!},$$

where the Stirling number  $\{k\}^n$  is the number of partitions into exactly  $k$  subsets.

Need more examples? If the coefficients of  $F(z)$  are arbitrary nonnegative numbers with a finite sum  $S$ , then  $F(z)/S$  defines a discrete probability distribution, and the convolution polynomial  $F_n(x)$  is  $S^x$  times the probability of obtaining the value  $n$  as the sum of  $x$  independent random variables having that distribution.

**A derived convolution.** Every convolution family  $\{F_n(x)\}$  satisfies another general convolution formula in addition to the one we began with:

$$(x + y) \sum_{k=0}^n k F_k(x) F_{n-k}(y) = x n F_n(x + y).$$

For example, if  $F_n(x)$  is the convolution family corresponding to powers of  $\mathcal{B}_t(z)$ , this formula says that

$$(x + y) \sum_{k=0}^n k \binom{x + tk}{k} \frac{x}{x + tk} \binom{y + t(n - k)}{n - k} \frac{y}{y + t(n - k)} = x n \binom{x + y + tn}{n} \frac{x + y}{x + y + tn};$$

it looks messy, but it simplifies to another amazing identity in four parameters,

$$\sum_{k=0}^n \binom{x + t(n - k)}{n - k} \binom{y + tk}{k} \frac{y}{y + tk} = \binom{x + y + tn}{n}$$

if we replace  $n$  by  $n + 1$ ,  $k$  by  $n + 1 - k$ , and  $x$  by  $x - t + 1$ . This identity has an interesting history going back to Rothe in 1793 see [Gould and Kaucký 1966].

The alternative convolution formula is proved by differentiating the basic identity  $F(z)^x = \sum_{n \geq 0} F_n(x) z^n$  with respect to  $z$  and multiplying by  $z$ :

$$x z F'(z) F(z)^{x-1} = \sum_{n \geq 0} n F_n(x) z^n.$$

Now  $\sum_{k=0}^n k F_k(x) F_{n-k}(y)$  is the coefficient of  $z^n$  in  $x z F'(z) F(z)^{x+y-1}$ , while  $n F_n(x + y)$  is the coefficient of  $z^n$  in  $(x + y) z F'(z) F(z)^{x+y-1}$ . Q.E.D.

**Convolution and composition.** Once upon a time I was trying to remember the form of a general convolution family, so I gave *Mathematica* the following command:

```
Simplify[Series[(1+Sum[A[k]z^k,{k,4}])^x,{z,0,4}]]
```

The result was a surprise. Instead of presenting the coefficient of  $z^n$  as a polynomial in  $x$ , *Mathematica* chose another form: The coefficient of  $z^2$ , for example, was  $\frac{1}{2}A_1^2x(x-1) + A_2x$ . In the notation of falling factorial powers, *Mathematica*'s response took the form

$$1 + A_1xz + \left(\frac{1}{2}A_1^2x^{\underline{2}} + A_2x\right)z^2 + \left(\frac{1}{6}A_1^3x^{\underline{3}} + A_1A_2x^{\underline{2}} + A_3x\right)z^3 \\ + \left(\frac{1}{24}A_1^4x^{\underline{4}} + \frac{1}{2}A_1^2A_2x^{\underline{3}} + \left(A_1A_3 + \frac{1}{2}A_2^2\right)x^{\underline{2}} + A_4x\right)z^4 + O(z)^5.$$

I wasn't prepared to work with factorial powers, so I tried another tack:

```
Simplify[Series[Exp[Sum[a[k]z^k,{k,4}]x],{z,0,4}]]
```

This time I got ordinary polynomials in  $x$ , but—lo and behold—they were

$$1 + a_1xz + \left(\frac{1}{2}a_1^2x^2 + a_2x\right)z^2 + \left(\frac{1}{6}a_1^3x^3 + a_1a_2x^2 + a_3x\right)z^3 \\ + \left(\frac{1}{24}a_1^4x^4 + \frac{1}{2}a_1^2a_2x^3 + \left(a_1a_3 + \frac{1}{2}a_2^2\right)x^2 + a_4x\right)z^4 + O(z)^5.$$

The result was exactly the same as before, but with  $a$ 's in place of  $A$ 's, and with normal powers in place of the factorials!

So I learned a curious phenomenon: *If we take any convolution family and replace each power  $x^k$  by  $x^{\underline{k}}$ , we get another convolution family.* (By the way, the replacement can be done in *Mathematica* by saying

```
Expand[F[n,x]]/.Power[x,k_]->k!Binomial[x,k];
```

expansion is necessary in case  $F_n(x)$  has been factored.)

The proof was not difficult to find, once I psyched out how *Mathematica* might have come up with its factorial-based formula: We have

$$e^{xf(z)} = 1 + f(z)x + \frac{f(z)^2}{2!}x^2 + \frac{f(z)^3}{3!}x^3 + \cdots,$$

and furthermore

$$(1 + f(z))^x = 1 + f(z)x + \frac{f(z)^2}{2!}x^{\underline{2}} + \frac{f(z)^3}{3!}x^{\underline{3}} + \cdots.$$

Therefore if we start with the convolution family  $F_n(x)$  corresponding to  $F(z) = e^{f(z)}$ , and replace each  $x^k$  by  $x^{\underline{k}}$ , we get the convolution family corresponding to  $1 + f(z) = 1 + \ln F(z)$ .

A similar derivation shows that if we replace  $x^k$  by the *rising* factorial power  $x^{\overline{k}}$  instead, we get the convolution family corresponding to  $1/(1 - f(z)) = 1/(1 - \ln F(z))$ . In particular, if we begin with the family  $F_n(x) = x(x+n)^{n-1}/n!$  corresponding to  $T(z)/z = e^{T(z)}$ , and if we replace  $x^k$  by  $x^{\overline{k}}$  to get

$$\frac{1}{n!} \sum_{k=0}^{n-1} \binom{n-1}{k} x^{\overline{k+1}} n^{n-1-k},$$

this must be  $[z^n](1 - T(z))^{-x} = t_n(x)/n!$ , the tree polynomial.

Indeed, we can replace each  $x^k$  by  $k! G_k(x)$ , where  $\{G_k(x)\}$  is any convolution family whatever! The previous examples,  $x^k$  and  $x^{\overline{k}}$ , are merely the special cases  $k! \binom{x}{k}$  and  $k! \binom{x+k-1}{k}$  corresponding to two of the simplest and most basic families we have considered. In general we get

$$1 + f(z) G_1(x) + \frac{f(z)^2}{2!} 2! G_2(x) + \frac{f(z)^3}{3!} 3! G_3(x) + \cdots ,$$

which is none other than  $G(f(z))^x = G(\ln F(z))^x$ .

For example,  $G_k(x) = \binom{x+2k}{k} \frac{x}{x+2k} = x(x+2k-1)^{\overline{k-1}}/k!$  is the family corresponding to  $\mathcal{B}_2(z)$ . If we know the family  $F_n(x)$  corresponding to  $e^{f(z)}$  we can replace  $x^k$  by  $x(x+2k-1)^{\overline{k-1}}$ , thereby obtaining the family that corresponds to  $\mathcal{B}_2(f(z)) = (1 + \sqrt{1 - 4f(z)})/2f(z)$ .

**Convolution matrices.** I knew that such remarkable facts must have been discovered before, although they were new to me at the time. And indeed, it was not difficult to find them in books, once I knew what to look for. (Special cases of general theorems are not always easy to recognize, because any particular formula is a special case of infinitely many generalizations, almost all of which are false.)

In the special case that each polynomial  $F_n(x)$  has degree exactly  $n$ , i.e., when  $f_1 \neq 0$ , the polynomials  $n! F_n(x)$  are said to be of *binomial type* [Mullin and Rota 1970]. An extensive theory of such polynomial sequences has been developed [Rota et al 1973] [Garsia 1973] [Roman and Rota 1978], based on the theory of linear operators, and the reader will find it quite interesting to compare the instructive treatment in those papers to the related but rather different directions explored in the present work. A comprehensive exposition of the operator approach appears in [Roman 1984]. Actually, Steffensen had defined a concept called *poweroids*, many years earlier [Steffensen 1941], and poweroids are almost exactly the same as sequences of binomial type; but Steffensen apparently did not realize that his poweroids satisfy the convolution property, which we can readily deduce (with hindsight) from equations (6) and (7) of his paper.

Eri Jabotinsky introduced a nice way to understand the phenomena of convolution polynomials, by considering the infinite matrix of coefficients  $f_{nk}$  [Jabotinsky 1947]. Let us recapitulate the notation that was introduced informally above:

$$\begin{aligned} e^{xf(z)} &= F(z)^x = 1 + F_1(x)z + F_2(x)z^2 + \cdots ; \\ F_n(x) &= (f_{n1}x + f_{n2}x^2 + \cdots + f_{nn}x^n)/n! ; \\ f(z) &= f_1z + f_2z^2/2! + f_3z^3/3! + \cdots . \end{aligned}$$

Then Jabotinsky's matrix  $F = (f_{nk})$  is a lower triangular matrix containing the coefficients of  $n! F_n(x)$  in the  $n$ th row. The first few rows are

$$\begin{array}{ccccccc} f_1 & & & & & & \\ f_2 & & f_1^2 & & & & \\ f_3 & & 3f_1f_2 & & f_1^3 & & \\ f_4 & & 4f_1f_3 + 3f_2^2 & & 6f_1^2f_2 & & f_1^4 , \end{array}$$

as we saw earlier. In general,

$$f_{nk} = \sum \frac{n!}{1!^{k_1} k_1! 2!^{k_2} k_2! 3!^{k_3} k_3! \dots} f_1^{k_1} f_2^{k_2} f_3^{k_3} \dots ,$$

summed over all  $k_1, k_2, k_3, \dots \geq 0$  with

$$k_1 + k_2 + k_3 + \dots = k, \quad k_1 + 2k_2 + 3k_3 + \dots = n.$$

(The summation is over all partitions of the integer  $n$  into  $k$  parts, where  $k_j$  of the parts are equal to  $j$ .) We will call such an array a *convolution matrix*.

If each original coefficient  $f_j$  is an integer, all entries of the corresponding convolution matrix will be integers, because the complicated quotient of factorials in the sum is an integer—it is the number of ways to partition a set of  $n$  elements into  $k$  subsets with exactly  $k_j$  of the subsets having size  $j$ . Given the first column we can compute the other columns from left to right and from top to bottom by using the recurrence

$$f_{nk} = \sum_{j=1}^{n-k+1} \binom{n-1}{j-1} f_j f_{(n-j)(k-1)}.$$

This recurrence is based on set partitions on which the element  $n$  occurs in a subset of size  $j$ : There are  $\binom{n-1}{j-1}$  ways to choose the other  $j-1$  elements of the subset, and the factor  $f_{(n-j)(k-1)}$  corresponds to partitioning the remaining  $n-j$  elements into  $k-1$  parts.

For example, if each  $f_j = 1$ , the convolution matrix begins

$$\begin{array}{cccccc} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 3 & 1 & & & \\ 1 & 7 & 6 & 1 & & \\ 1 & 15 & 25 & 10 & 1 & \end{array}.$$

These are the numbers  $\{n_k\}$  that *Mathematica* calls **StirlingS2[n,k]**; they arose in our example of Bell numbers when  $f(z) = e^z - 1$ . Similarly, if each  $f_j = (j-1)!$ , the first five rows are

$$\begin{array}{cccccc} 1 & & & & & \\ 1 & 1 & & & & \\ 2 & 3 & 1 & & & \\ 6 & 11 & 6 & 1 & & \\ 24 & 50 & 35 & 10 & 1 & \end{array};$$

*Mathematica* calls these numbers  $(-1)^{n-k} \text{StirlingS1}[n,k]$ . In this case  $f(z) = \ln(1/(1-z))$ , and  $F_n(z) = \binom{x+n-1}{n}$ . The signed numbers **StirlingS1[nk]**,

$$\begin{array}{cccccc} 1 & & & & & \\ -1 & 1 & & & & \\ 2 & -3 & 1 & & & \\ -6 & 11 & -6 & 1 & & \\ 24 & -50 & 35 & -10 & 1 & \end{array}$$



correspond to  $f(z) = \ln(1+z)$  and  $F_n(z) = \binom{x}{n}$ . In general if we replace  $z$  by  $\alpha z$  and  $x$  by  $\beta x$ , the effect is to multiply row  $n$  of the matrix by  $\alpha^n$  and to multiply column  $k$  by  $\beta^k$ . Thus when  $\beta = \alpha^{-1}$ , the net effect is to multiply  $f_{nk}$  by  $\alpha^{n-k}$ . Transforming the signs by a factor  $(-1)^{n-k}$  corresponds to changing  $F(z)$  to  $1/F(-z)$  and  $f(z)$  to  $-f(-z)$ . Therefore the matrix that begins

$$\begin{array}{cccccc} 1 & & & & & \\ -1 & 1 & & & & \\ 1 & -3 & 1 & & & \\ -1 & 7 & -6 & 1 & & \\ 1 & -15 & 25 & -10 & 1 & \end{array}$$

corresponds to  $f(z) = 1 - e^{-z}$ .

Let's look briefly at some of our other examples in matrix form. When  $F(z) = \mathcal{B}_t(z)$ , we have  $f_j = (tj - 1)\frac{j-1}{2}$ , which is an integer when  $t$  is an integer. In particular, the Catalan case  $t = 2$  produces a matrix that begins

$$\begin{array}{cccccc} 1 & & & & & \\ 3 & 1 & & & & \\ 20 & 9 & 1 & & & \\ 210 & 107 & 18 & 1 & & \\ 3024 & 1650 & 335 & 30 & 1 & \end{array}$$

When  $t = 1/2$ , we can remain in an all-integer realm by replacing  $z$  by  $2z$  and  $x$  by  $x/2$ . Then  $f_j = 0$  when  $j$  is even, while  $f_{2j+1} = (-1)^j(2j-1)!!^2$ :

$$\begin{array}{cccccc} 1 & & & & & \\ 0 & 1 & & & & \\ -1 & 0 & 1 & & & \\ 0 & -4 & 0 & 1 & & \\ 9 & 0 & -10 & 0 & 1 & \end{array};$$

If we now replace  $z$  by  $iz$  and  $x$  by  $x/i$  to eliminate the minus signs, we find that  $f(z) = \arcsin z$ , because  $\ln(iz + \sqrt{1-z^2}) = i\theta$  when  $z = \sin \theta$ . Thus we can deduce a closed form for the coefficients of  $e^{x \arcsin z} = \mathcal{B}_{1/2}(2iz)^{x/(2i)}$ :

$$\begin{aligned} n! [z^n] e^{x \arcsin z} &= (2i)^{n-1} x \left( \frac{x}{2i} + \frac{n}{2} - 1 \right) \dots \left( \frac{x}{2i} - \frac{n}{2} + 1 \right) \\ &= \begin{cases} x^2(x^2 + 2^2) \dots (x^2 + (n-2)^2), & n \text{ even;} \\ x(x^2 + 1^2)(x^2 + 3^2) \dots (x^2 + ((n-2)^2)), & n \text{ odd.} \end{cases} \end{aligned}$$

This remarkable formula is equivalent to the theorem of [Gomes Teixeira 1896].

If  $f_j = 2^{1-j}$  when  $j$  is odd but  $f_j = 0$  when  $j$  is even, we get the convolution matrix corresponding to  $e^{2x \sinh(z/2)}$ :

$$\begin{array}{cccccc} 1 & & & & & \\ 0 & 1 & & & & \\ \frac{1}{4} & 0 & 1 & & & \\ 0 & 1 & 0 & 1 & & \\ \frac{1}{16} & 0 & \frac{5}{2} & 0 & 1 & \\ 0 & 1 & 0 & 5 & 0 & 1 \\ \frac{1}{64} & 0 & \frac{91}{16} & 0 & \frac{35}{4} & 0 & 1 \\ 0 & 1 & 0 & 21 & 0 & 14 & 0 & 1 \end{array}.$$

Again we could stay in an all-integer realm if we replaced  $z$  by  $2z$  and  $x$  by  $x/2$ ; but the surprising thing in this case is that the entries in even-numbered rows and columns are all integers *before* we make any such replacement. The reason is that the entries satisfy  $f_{nk} = k^2 f_{(n-2)k}/4 + f_{(n-2)(k-2)}$ . (See [Riordan 1968, pages 213–217], where the notation  $T(n, k)$  is used for these “central factorial numbers”  $f_{nk}$ .)

We can complete our listing of noteworthy examples by setting  $f_j = \sum_{k=1}^n n^{n-k-1} n^{\underline{k}}$ ; then we get the coefficients of the tree polynomials:

$$\begin{array}{ccccccc} & & 1 & & & & \\ & 3 & & 1 & & & \\ & 17 & & 9 & & 1 & \\ & 142 & & 95 & & 18 & & 1 \\ 1569 & 1220 & 305 & 30 & 1 & & . \end{array}$$

The sum of the entries in row  $n$  is  $n^n$ .

**Composition and iteration.** Jabotinski’s main reason for defining things as he did was his observation that *the product of convolution matrices is a convolution matrix*. Indeed, if  $F$  and  $G$  are the convolution matrices corresponding to the functions  $e^{xf(z)}$  and  $e^{xg(z)}$  we have the vector/matrix identities

$$\begin{aligned} e^{xf(z)} - 1 &= (z, z^2/2!, z^3/3!, \dots) F(x, x^2, x^3, \dots)^T \\ e^{xg(z)} - 1 &= (z, z^2/2!, z^3/3!, \dots) G(x, x^2, x^3, \dots)^T \end{aligned}$$

If we now replace  $x^k$  in  $e^{xf(z)}$  by  $k! G_k(x)$ , as in our earlier discussion, we get

$$\begin{aligned} &(z, z^2/2!, z^3/3!, \dots) F(G_1(x), 2! G(x), 3! G_3(x), \dots)^T \\ &= (z, z^2/2!, z^3/3!, \dots) FG(x, x^2, x^3, \dots)^T \\ &= (f(z), f(z)^2/2!, f(z)^3/3!, \dots) G(x, x^2, x^3, \dots)^T \\ &= e^{xg(f(z))} - 1. \end{aligned}$$

Multiplication of convolution matrices corresponds to composition of the functions in the exponent.

Why did the function corresponding to  $FG$  turn out to be  $g(f(z))$  instead of  $f(g(z))$ ? Jabotinsky, in fact, defined his matrices as the transposes of those given here. The rows of his (upper triangular) matrices were the power series  $f(z)^k$ , while the columns were the polynomials  $F_n(x) = [z^n] e^{xf(z)}$ ; with those conventions the product of his matrices  $F^T G^T$  corresponded to  $f(g(z))$ . (In fact, he defined a considerably more general representation, in which the matrix  $F$  could be  $U^{-1} F U$  for any nonsingular matrix  $U$ .) However, when our interest is focussed on the polynomials  $n! F_n(x)$ , as when we study Stirling numbers or tree polynomials or the Stirling polynomials to be discussed below, it is more natural to work with lower triangular matrices and to insert factorial coefficients, as Comtet did [Comtet 1970, section 3.7]. The two conventions are isomorphic. Without the factorials, convolution matrices are sometimes called *renewal arrays* [Rogers 1978]. We would get a non-reversed order if we had been accustomed to using postfix notation

$(z)f$  for functions, as we do for operations such as squaring or taking transposes or factorials; then  $g(f(z))$  would be  $((z)f)g$ .

Recall that the Stirling numbers  $\{n_k\}$  correspond to  $f(z) = e^z - 1$ , and the Stirling numbers  $[n_k]$  correspond to  $g(z) = \ln(1/(1-z))$ . Therefore if we multiply Stirling's triangles we get the convolution matrix

$$FG = \begin{array}{ccccc} & 1 & & & \\ & 2 & 1 & & \\ & 6 & 6 & 1 & \\ & 26 & 36 & 12 & 1 \\ 150 & 250 & 120 & 20 & 1 \end{array},$$

which corresponds to  $g(f(z)) = \ln(1/(2 - e^z))$ . Voila! These convolution polynomials represent the coefficients of  $(2 - e^z)^{-x}$ . [Cayley 1859] showed that  $(2 - e^z)^{-1}$  is the exponential generating function for the sequence 1, 3, 13, 75, 541,  $\dots$ , which counts *preferential arrangements* of  $n$  objects, i.e., different outcomes of sorting when equality is possible as well as inequality. The coefficient  $(fg)_{nk}$  is the number of preferential arrangements in which the “current minimum” changes  $k$  times when we examine the elements one by one in some fixed order. (See [Graham et al 1989, exercise 7.44].)

Similarly, the reverse matrix product yields the so-called Lah numbers [Lah 1955],

$$GF = \begin{array}{ccccc} & 1 & & & \\ & 2 & 1 & & \\ & 6 & 6 & 1 & \\ & 24 & 36 & 12 & 1 \\ 120 & 240 & 120 & 20 & 1 \end{array};$$

here  $f_j = j!$  and the rows represent the coefficients of  $\exp(xf(g(z))) = \exp(xz + xz^2 + xz^3 + \dots)$ . Indeed, the convolution polynomials in this case are the generalized Laguerre polynomials  $L_n^{(-1)}(-x)$ , which *Mathematica* calls `LaguerreL[n,-1,-x]`. These polynomials can also be expressed as  $L_n(-x) - L_{n-1}(-x)$ ; or as `LaguerreL[n,-x]-LaguerreL[n-1,-x]` if we say

```
Unprotect[LaguerreL]; LaguerreL[-1,x_]:=0; Protect[LaguerreL]
```

first. The row sums 1, 3, 13, 73, 501,  $\dots$  of  $GF$  enumerate “sets of lists” [Motzkin 1971]; the coefficients are  $(GF)_{nk} = n! [z^n] f(g(z))^k / k! = \binom{n}{k} \binom{n-1}{k-1} (n-k)!$  [Riordan 1968, exercise 5.7].

Since convolution matrices are closed under multiplication, they are also closed under exponentiation, i.e., under taking of powers. The  $q$ th power  $F^q$  of a convolution matrix then corresponds to  $q$ -fold iteration of the function  $\ln F = f$ . Let us denote  $f(f(z))$  by  $f^{[2]}(z)$ ; in general, the  $q$ th iterate  $f^{[q]}(z)$  is defined to be  $f(f^{[q-1]}(z))$ , where  $f^{[0]}(z) = z$ . This is *Mathematica*'s `Nest[f,z,q]`.

The  $q$ th iterate can be obtained by doing  $O(\log q)$  matrix multiplications, but in the interesting case  $f'(0) = f_1 = 1$  we can also compute the coefficients of  $f^{[q]}(z)$  by using formulas in which  $q$  is simply a numerical parameter. Namely, as suggested by [Jabotinsky 1947], we can express the matrix power  $F^q$  as

$$(I + (F - I))^q = I + \binom{q}{1}(F - I) + \binom{q}{2}(F - I)^2 + \binom{q}{3}(F - I)^3 + \dots$$

This infinite series converges, because the entry in row  $n$  and column  $k$  of  $(F - I)^j$  is zero for all  $j > n - k$ . When  $q$  is any positive integer, the result defined in this way is a convolution matrix. Furthermore, the matrix entries are all polynomials in  $q$ . Therefore the matrix obtained by this infinite series is a convolution matrix for all values of  $q$ .

Another formula for the entries of  $F^q$  was presented in [Jabotinsky 1963]. Let  $f_{nk}^{(q)}$  be the element in row  $n$  and column  $k$ ; then

$$\begin{aligned} f_{nk}^{(q)} &= \sum_{l=0}^m \binom{q}{l} (F - I)_{nk}^l \\ &= \sum_{j=0}^m f_{nk}^{(j)} \sum_{l=j}^m \binom{q}{l} \binom{l}{j} (-1)^{l-j} \\ &= \sum_{j=0}^m f_{nk}^{(j)} \binom{q}{j} \sum_{l=j}^m \binom{q-j}{l-j} (-1)^{l-j} \\ &= \sum_{j=0}^m f_{nk}^{(j)} \binom{q}{j} \binom{q-j-1}{m-j} (-1)^{m-j}, \end{aligned}$$

for any  $m \geq n - k$ . Indeed, we have  $p(q) = \sum_{j=0}^m p(j) \binom{q}{j} \binom{q-j-1}{m-j} (-1)^{m-j}$  whenever  $p$  is a polynomial of degree  $\leq m$ ; this is a special case of Lagrange interpolation.

It is interesting to set  $q = 1/2$  and compute convolution square roots of the Stirling number matrices. We have

$$\begin{aligned} \begin{pmatrix} 1 & & & & \\ 1/2 & 1 & & & \\ 1/8 & 3/2 & 1 & & \\ 0 & 5/4 & 3 & 1 & \\ 1/32 & 5/8 & 5 & 5 & 1 \end{pmatrix}^2 &= \begin{pmatrix} 1 & & & & \\ 1 & 1 & & & \\ 1 & 3 & 1 & & \\ 1 & 7 & 6 & 1 & \\ 1 & 15 & 25 & 10 & 1 \end{pmatrix}; \\ \begin{pmatrix} 1 & & & & \\ 1/2 & 1 & & & \\ 5/8 & 3/2 & 1 & & \\ 5/4 & 13/4 & 3 & 1 & \\ 109/32 & 75/8 & 10 & 5 & 1 \end{pmatrix}^2 &= \begin{pmatrix} 1 & & & & \\ 1 & 1 & & & \\ 2 & 3 & 1 & & \\ 6 & 11 & 6 & 1 & \\ 24 & 50 & 35 & 10 & 1 \end{pmatrix}. \end{aligned}$$

The function  $z + z^2/4 + z^3/48 + z^5/3840 - 7z^6/92160 + \dots$  therefore lies “halfway” between  $z$  and  $e^z - 1 = z + z^2/2! + z^3/3! + \dots$ , and the function  $z + z^2/4 + 5z^3/48 + 5z^4/96 + 109z^5/3840 + 497z^6/30720 + \dots$  lies halfway between  $z$  and  $\ln 1/(1 - z) = z + z^2/2 + z^3/3 + \dots$ . These half-iterates are unfamiliar functions; but it is not difficult to prove that  $z/(1 - z/2) = z + z^2/2 + z^3/4 + \dots$  is halfway between  $z$  and  $z/(1 - z) = z + z^2 + z^3 + \dots$ . In general when  $f(z) = z/(1 - cz^k)^{1/k}$  we have  $f^{[q]}(z) = z/(1 - qc^k z^k)^{1/k}$ .

It seems natural to conjecture that the coefficients of  $f^{[q]}(z)$  are positive for  $q > 0$  when  $f(z) = \ln 1/(1 - z)$ ; but this conjecture turns out to be false, because *Mathematica* reports that

$[z^8] f^{[q]}(z) = -11q/241920 + O(q^2)$ . Is there a simple necessary and sufficient condition on  $f$  that characterizes when all coefficients of  $f^{[q]}$  are nonnegative for nonnegative  $q$ ? This will happen if and only if the entries in the first column of

$$\ln F = (F - I) - \frac{1}{2}(F - I)^2 + \frac{1}{3}(F - I)^3 - \dots$$

are nonnegative. (See [Kuczma 1968] for iteration theory and an extensive bibliography.)

**Reversion.** The case  $q = -1$  of iteration is often called reversion of series, although *Mathematica* uses the more proper name **InverseSeries**. Given  $f(z) = f_1 z + f_2 z^2/2! + \dots$ , we seek  $g(z) = f^{[-1]}(z)$  such that  $g(f(z)) = z$ . This is clearly equivalent to finding the first column of the inverse of the convolution matrix.

The inverse does not exist when  $f_1 = 0$ , because the diagonal of  $F$  is zero in that case. Otherwise we can assume that  $f_1 = 1$ , because  $f_1 g(f(z)/f_1) = z$  when  $g$  reverts the power series  $f(z/f_1)$ .

When  $f_1 = 1$  we can obtain the inverse by setting  $q = -1$  in our general formula for iteration. But Lagrange's celebrated inversion theorem for power series tells us that there is another, more informative, way to compute the function  $g = f^{[-1]}$ . Let us set  $\hat{F}(z) = f(z)/z = 1 + f_2 z/2! + f_3 z^2/3! + \dots$ . Then Lagrange's theorem states that the elements of the matrix  $G = F^{-1}$  are

$$g_{nk} = \frac{(n-1)!}{(k-1)!} \hat{F}_{n-k}(-n),$$

where  $\hat{F}_n(x)$  denotes the convolution family corresponding to  $\hat{F}(z)$ .

There is a surprisingly simple way to prove Lagrange's theorem, using our knowledge of convolution families. Note first that

$$f_{nk} = n! [z^n x^k] e^{xf(z)} = \frac{n!}{k!} [z^n] f(z)^k = \frac{n!}{k!} [z^{n-k}] \hat{F}(z)^k;$$

therefore

$$f_{nk} = \frac{n!}{k!} \hat{F}_{n-k}(k).$$

Now we need only verify that the matrix product  $GF$  is the identity, by computing its element in row  $n$  and column  $m$ :

$$\sum_{k=m}^n g_{nk} f_{km} = \sum_{k=m}^n \frac{(n-1)!}{(k-1)!} \hat{F}_{n-k}(-n) \frac{k!}{m!} \hat{F}_{k-m}(m).$$

When  $m = n$  the sum is obviously 1. When  $m = n - p$  for  $p > 0$  it is  $(n-1)!/(n-p)!$  times

$$\begin{aligned} \sum_{k=n-p}^n k \hat{F}_{n-k}(-n) \hat{F}_{k-n+p}(n-p) &= \sum_{k=0}^p (n-k) \hat{F}_k(-n) \hat{F}_{p-k}(n-p) \\ &= n \sum_{k=0}^p \hat{F}_k(-n) \hat{F}_{p-k}(n-p) - \sum_{k=0}^p k \hat{F}_k(-n) \hat{F}_{p-k}(n-p) \\ &= n \hat{F}_p(-p) - n \hat{F}_p(-p) = 0 \end{aligned}$$

by the original convolution formula and the one we derived from it. The proof is complete.

**Extending the matrix.** The simple formula for  $f_{nk}$  that we used to prove Lagrange's theorem when  $f_1 = 1$  can be written in another suggestive form, if we replace  $k$  by  $n - k$ :

$$f_{n(n-k)} = n^k \widehat{F}_k(n - k).$$

For every fixed  $k$ , this is a polynomial in  $n$ , of degree  $\leq 2k$ . Therefore we can define the quantity  $f_{y(y-k)}$  for all real or complex  $y$  to be  $y^k \widehat{F}_k(y - k)$ ; and in particular we can define  $f_{nk}$  in this manner for all integers  $n$  and  $k$ , letting  $f_{nk} = 0$  when  $k > n$ . For example, in the case of Stirling numbers this analysis establishes the well-known fact that  $\left\{ \begin{smallmatrix} y \\ y-k \end{smallmatrix} \right\}$  and  $\left[ \begin{smallmatrix} y \\ y-k \end{smallmatrix} \right]$  are polynomials in  $y$  of degree  $2k$ , and that these polynomials are multiples of  $y^{\overline{k+1}} = y(y-1) \dots (y-k)$  when  $k > 0$ .

The two flavors of Stirling numbers are related in two important ways. First, their matrices are inverse to each other if we attach the signs  $(-1)^{n-k}$  to the elements in one matrix:

$$\sum_{k=0}^n \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} \left[ \begin{smallmatrix} k \\ m \end{smallmatrix} \right] (-1)^{n-k} = \sum_{k=0}^m \left[ \begin{smallmatrix} n \\ k \end{smallmatrix} \right] \left\{ \begin{smallmatrix} k \\ m \end{smallmatrix} \right\} (-1)^{n-k} = \delta_{mn}.$$

This follows since the numbers  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$  correspond to  $f(z) = e^z - 1$  and the numbers  $\left[ \begin{smallmatrix} n \\ k \end{smallmatrix} \right] (-1)^{n-k}$  correspond to  $g(z) = \ln(1+z)$ , as mentioned earlier, and we have  $g(f(z)) = z$ .

The other important relationship between  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$  and  $\left[ \begin{smallmatrix} n \\ k \end{smallmatrix} \right]$  is the striking identity

$$\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} = \left[ \begin{smallmatrix} -k \\ -n \end{smallmatrix} \right],$$

which holds for all integers  $n$  and  $k$  when we use the polynomial extension method. We can prove in fact, that the analogous relation

$$f_{nk} = (-1)^{k-n} g_{(-k)(-n)}$$

holds in the extended matrices  $F$  and  $G$  that correspond to *any* pair of inverse functions  $g(f(z)) = z$ , when  $f'(0) = 1$ . For we have

$$(-1)^{n-k} g_{(-k)(-n)} = (-1)^{n-k} (-k-1)(-k-2) \dots (-n) \widehat{F}_{n-k}(k) = \frac{n!}{k!} \widehat{F}_{n-k}(k) = f_{nk}$$

in the formulas above. (The interesting history of the identity  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} = \left[ \begin{smallmatrix} -k \\ -n \end{smallmatrix} \right]$  is traced in [Knuth 1992]. The fact that the analogous formula holds in any convolution matrix was pointed out by Ira Gessel after he had read a draft of that paper. See also [Jabotinski 1953]; [Carlitz 1978]; [Roman and Rota 1978, section 10].)

Suppose we denote the Lah numbers  $\binom{n}{k} \binom{n-1}{k-1} (n-k)!$  by  $\left| \begin{smallmatrix} n \\ k \end{smallmatrix} \right|$ . The extended matrix in that case has a pleasantly symmetrical property

$$\left| \begin{smallmatrix} n \\ k \end{smallmatrix} \right| = \left| \begin{smallmatrix} -k \\ -n \end{smallmatrix} \right|,$$

because the corresponding function  $f(z) = z/(1 - z)$  satisfies  $f(-f(-z)) = z$ . (Compare [Mullin and Rota 1969, section 9].) Near the origin  $n = k = 0$ , the nonzero entries look like this:

$$\begin{array}{ccccccc}
\dots & 1 & & & & & \\
\dots & 12 & 1 & & & & \\
\dots & 36 & 6 & 1 & & & \\
\dots & 24 & 6 & 2 & 1 & & \\
& & & & 1 & & \\
& & & & & 1 & \\
& & & & & 2 & 1 \\
& & & & & 6 & 6 & 1 \\
& & & & & 24 & 36 & 12 & 1 \\
& & & & & \vdots & \vdots & \vdots & \\
& & & & & & & & 
\end{array}$$

**Still more convolutions.** Our proof of Lagrange's theorem yields yet another corollary. Suppose  $g(f(z)) = z$  and  $f'(0) = 1$ , and let  $\widehat{F}(z) = f(z)/z$ ,  $\widehat{G}(z) = g(z)/z$ . Then the equation

$$g_{nk} = \frac{n!}{k!} \hat{G}_{n-k}(k) = \frac{(n-1)!}{(k-1)!} \hat{F}_{n-k}(-n)$$

tell us, after replacing  $n$  by  $n + k$ , that the identity

$$\frac{n+k}{k} \hat{G}_n(k) = \hat{F}_n(-n-k)$$

holds for all positive integers  $k$ . Thus the polynomials  $\widehat{G}_n(x)$  and  $\widehat{F}_n(x)$  must be related by the formula

$$(x+n)\hat{G}_n(x) = x\hat{F}_n(-x-n).$$

Now  $\widehat{F}_n(x)$  is an arbitrary convolution family, and  $\widehat{F}_n(-x)$  is another. We can conclude that if  $\{F_n(x)\}$  is any convolution family, then so is the set of polynomials  $\{xF_n(x+n)/(x+n)\}$ . Indeed, if  $F_n(x)$  corresponds to the coefficients of  $F(z)^x$ , our argument proves that the coefficients of  $G(z)^x$  are  $x F_n(x+n)/(x+n)$ , where  $zG(z)$  is the inverse of the power series  $z/F(z)$ :

$$G(z) = F(zG(z)) \,, \quad G(z/F(z)) = F(z) \,.$$

The case  $F(z) = 1 + z$  and  $G(z) = 1/(1 - z)$  provides a simple example, where we know that  $F_n(x) = \binom{x}{n}$  and  $G_n(x) = \binom{x+n-1}{n} = xF_n(x+n)/(x+n)$ .

A more interesting example arises when  $F(z) = ze^z/(e^z - 1) = z + z/(e^z - 1) = 1 + z/2 + B_2 z^2/2! + B_4 z^4/4! + \dots$ ; then  $F(-z)$  is the exponential generating function for the Bernoulli numbers. The convolution family for  $F(z)^x$  is  $F_n(x) = x\sigma_n(x)$ , where  $\sigma_n(x)$  is called a *Stirling polynomial*. (Actually  $\sigma_0(x) = 1/x$ , but  $\sigma_n(x)$  is a genuine polynomial when  $n \geq 1$ .) The function  $G$  such that  $G(z/F(z)) = F(z)$  is  $G(z) = z^{-1} \ln(1/(1-z))$ ; therefore the convolution family for  $G(z)^x$  is  $G_n(x) = xF_n(x+n)/(x+n) = x\sigma_n(x+n)$ .

In this example the convolution family for  $e^{xzG(z)} = (1-z)^{-x}$  is

$$\binom{x+n-1}{n} = \frac{1}{n!} \left( \binom{n}{0} + \binom{n}{1}x + \cdots + \binom{n}{n}x^n \right);$$

therefore

$$\left[ \begin{matrix} n \\ n-k \end{matrix} \right] = \frac{n!}{(n-k)!} G_k(n-k) = \frac{n!}{(n-k)!} (n-k) \sigma_k(n) = n(n-1) \cdots (n-k) \sigma_k(n).$$

We also have

$$\left\{ \begin{matrix} n \\ n-k \end{matrix} \right\} = \left[ \begin{matrix} k-n \\ -n \end{matrix} \right] = (k-n)(k-1-n) \cdots (-n) \sigma_k(k-n).$$

These formulas, which are polynomials in  $n$  of degree  $2k$  for every fixed  $k$ , explain why the  $\sigma$  functions are called Stirling polynomials. Notice that  $\sigma_n(1) = (-1)^n B_n/n!$ ; it can also be shown that  $\sigma_n(0) = -B_n/(n \cdot n!)$ .

The process of going from  $F_n(x)$  to  $xF_n(x+n)/(x+n)$  can be iterated: Another replacement gives  $xF_n(x+2n)/(x+2n)$ , and after  $t$  iterations we discover that the polynomials  $xF_n(x+tn)/(x+tn)$  also form a convolution family. This holds for all positive integers  $t$ , and the convolution condition is expressible as a set of polynomial relations in  $t$ ; *therefore  $xF_n(x+tn)/(x+tn)$  is a convolution family for all complex numbers  $t$* . If  $F_n(x) = [z^n] F(z)^x$ , then  $xF_n(x+tn)/(x+tn) = [z^n] \mathcal{F}_t(z)^x$ , where  $\mathcal{F}_t(z)$  is defined implicitly by the equation

$$\mathcal{F}_t(z) = F(z\mathcal{F}_t(z)^t).$$

In particular, we could have deduced the convolution properties of the coefficients of  $\mathcal{B}_t(z)^x$  in this way.

Let us restate what we have just proved, combining it with the “derived convolution formula” obtained earlier:

**Theorem.** *Let  $F_n(x)$  be any family of polynomials in  $x$  such that  $F_n(x)$  has degree  $\leq n$ . If*

$$F_n(2x) = \sum_{k=0}^n F_k(x) F_{n-k}(x)$$

*holds for all  $n$  and  $x$ , then the following identities hold for all  $n$ ,  $x$ ,  $y$ , and  $t$ :*

$$\begin{aligned} \frac{(x+y)F_n(x+y+tn)}{x+y+tn} &= \sum_{k=0}^n \frac{x F_k(x+tk)}{x+tk} \frac{y F_{n-k}(y+t(n-k))}{y+t(n-k)}; \\ \frac{n F_n(x+y+tn)}{x+y+tn} &= \sum_{k=1}^n \frac{k F_k(x+tk)}{x+tk} \frac{y F_{n-k}(y+t(n-k))}{y+t(n-k)}. \end{aligned}$$

**Additional constructions.** We have considered several ways to create new convolution families from given ones, by multiplication or exponentiation of the associated convolution matrices, or by



replacing  $F_n(x)$  by  $x F_n(x + tn)/(x + tn)$ . It is also clear that the polynomials  $\alpha^n F_n(\beta x)$  form a convolution family whenever the polynomials  $F_n(x)$  do.

One further operation deserves to be mentioned: If  $F_n(x)$  and  $G_n(x)$  are convolution families, then so is the family  $H_n(x)$  defined by

$$H_n(x) = \sum_{k=0}^n F_k(x) G_{n-k}(x).$$

This is obvious, since  $H_n(x) = [z^n] F(z)^x G(z)^x$ . The corresponding operation on matrices  $F = (f_{nk})$ ,  $G = (g_{nk})$ ,  $H = (h_{nk})$  is

$$h_{nk} = \sum_{i,j} \binom{n}{j} f_{ji} g_{(n-j)(k-i)}.$$

If we denote this binary operation by  $H = F \circ G$ , it is interesting to observe that the associative law holds:  $(E \circ F) \circ G = E \circ (F \circ G)$  is true for all matrices  $E$ ,  $F$ ,  $G$ , not just for convolution matrices. A convolution matrix is characterized by the special property  $F \circ F = F \operatorname{diag}(2, 4, 8, \dots)$ .

The construction just mentioned is merely a special case of the one-parameter family

$$H_n^{(t)}(x) = \sum_{k=0}^n F_k(x) G_{n-k}(x + tk).$$

Again,  $\{H_n^{(t)}(x)\}$  turns out to be a convolution family, for arbitrary  $t$ : We have

$$\begin{aligned} \sum_{k=0}^n H_n^{(t)}(x) z^n &= \sum_{n \geq k \geq 0} F_k(x) G_{n-k}(x + tk) z^n = \sum_{n, k \geq 0} F_k(x) G_n(x + tk) z^{n+k} \\ &= \sum_{k \geq 0} F_k(x) z^k G(z)^{x+tk} = G(z)^x F(z G(z)^t)^x, \end{aligned}$$

so  $H_n(x) = [z^n] (G(z) F(z G(z)^t))^x$ .

**Applications.** What's the use of all this? Well, we have shown that many interesting convolution families exist, and that we can deduce nonobvious facts with comparatively little effort once we know that we're dealing with a convolution family.

One moral to be drawn is therefore the following. Whenever you encounter a triangular pattern of numbers that you haven't seen before, check to see if the first three rows have the form

$$\begin{array}{ccc} a & & \\ b & a^2 & \\ c & 3ab & a^3 \end{array}$$

for some  $a, b, c$ . (You may have to multiply or divide the  $n$ th row by  $n!$  first, and/or reflect its entries left to right.) If so, and if the problem you are investigating is mathematically "clean," chances are good that the fourth row will look like

$$d \quad 4ac + 3b^2 \quad 6a^2b \quad a^4.$$

And if so, chances are excellent that you are dealing with a convolution family. And if so, you may well be able to solve your problem.

In fact, exactly that scenario has helped the author on several occasions.

**Asymptotics.** Once you have identified a convolution family  $F_n(x)$ , you may well want to know the approximate value of  $F_n(x)$  when  $n$  and  $x$  are large. The remainder of this paper discusses a remarkable general power series expansion, discovered with the help of *Mathematica*, which accounts for the behavior of  $F_n(x)$  when  $n/x$  stays bounded and reasonably small as  $x \rightarrow \infty$ , although  $n$  may also vary as a function of  $x$ . We will assume that  $F_n(x)$  is the coefficient of  $z^n$  in  $F(z)^x$ , where  $F(0) = F'(0) = 1$ .

Our starting point is the classical “saddle point method,” which shows that in many cases the coefficient of  $z^n$  in a power series  $P(z)$  can be approximated by considering the value of  $P$  at a point where the derivative of  $P(z)/z^n$  is zero. (See [Good 1957].) In our case we have  $P(z) = e^{xf(z)}$ , where  $f(z) = \ln F(z) = z + f_2 z^2/2! + \dots$ ; and the derivative is zero when  $xf'(z) = n/z$ . Let this saddle point occur at  $z = s$ ; thus, we have

$$s f'(s) = n/x.$$

Near  $s$  we have  $f(z) = f(s) + (z - s)f'(s) + O((z - s)^2)$ ; so we will base our approximation on the assumption that the  $O((z - s)^2)$  contribution is zero. The approximation to  $F_n(x)$  will be  $\tilde{F}_n(x)$ , where

$$\begin{aligned} \tilde{F}_n(x) &= [z^n] \exp(x f(s) + x(z - s) f'(s)) \\ &= \frac{e^{x(f(s) - s f'(s))}}{n!} x^n f'(s)^n = \frac{F(s)^x}{n!} \left(\frac{n}{es}\right)^n. \end{aligned}$$

First let's look at some examples; later we will show that the ratio  $F_n(x)/\tilde{F}_n(x)$  is well behaved as a formal power series. Throughout this discussion we will let

$$y = n/x;$$

our goal, remember, is to find approximations that are valid when  $y$  is not too large, as  $x$  and possibly  $n$  go to  $\infty$ .

The simplest example is, of course,  $F(z) = e^z$  and  $f(z) = z$ ; but we needn't sneeze at it because it will give us some useful calibration. In this case  $f''(z) = 0$ , so our approximation will be exact. We have  $s = y$ , hence

$$\tilde{F}_n(x) = \frac{e^{xy}}{n!} \left(\frac{n}{ey}\right)^n = \frac{e^n}{n!} \left(\frac{x}{e}\right)^n = \frac{x^n}{n!} = F_n(x).$$

Next let's consider the case  $F(z) = T(z)/z$ ,  $f(z) = T(z)$ , when we know that  $F_n(x) = x(x + n)^{n-1}/n!$ . In this case  $zT'(z) = T(z)/(1 - T(z))$ , so we have  $T(s)/(1 - T(s)) = y$  or

$$T(s) = \frac{y}{1 + y}, \quad s = \frac{y}{1 + y} e^{-y/(1+y)}$$

because  $T(z) = ze^{T(z)}$ . Therefore

$$\tilde{F}_n(x) = \frac{e^{xy/(1+y)}}{n!} \left( \frac{n(1+y)}{ey e^{-y/(1+y)}} \right)^n = \frac{(x+n)^n}{n!};$$

the ratio  $F_n(x)/\tilde{F}_n(x) = x/(x+n) = 1/(1+y)$  is indeed near 1 when  $y$  is small.

If  $F(z) = 1+z$  we find, similarly,  $s = y/(1-y)$  and

$$n! \tilde{F}_n(x) = \left( \frac{1}{1-y} \right)^x \left( \frac{n(1-y)}{ey} \right)^n = \frac{x^x e^{-n}}{(x-n)^{x-n}};$$

by Stirling's approximation we also have

$$n! F_n(x) = \frac{x!}{(x-n)!} = \frac{x^x e^{-n}}{(x-n)^{x-n}} (1-y)^{-1/2} (1 + O(x^{-1})).$$

Again the ratio  $F_n(x)/\tilde{F}_n(x)$  is near 1. In general if  $F(z) = \mathcal{B}_t(z)$  the saddle point  $s$  turns out to be  $y(1+(t-1)y)^{t-1}/(1+ty)^t$ , and

$$n! \tilde{F}_n(x) = \frac{(x+tn)^{x+tn} e^{-n}}{(x+(t-1)n)^{x+(t-1)n}};$$

a similar analysis shows that this approximation is quite good, for any fixed  $t$ .

We know that

$$F_n(x) = \frac{x^n}{n!} \left( 1 + \frac{f_{n(n-1)}}{x} + \frac{f_{n(n-2)}}{x^2} + \dots \right)$$

and that  $f_{n(n-k)}$  is always a polynomial in  $n$  of degree  $\leq 2k$ . Therefore if  $n^2/x \rightarrow 0$  as  $x \rightarrow \infty$ , we can simply use the approximation  $F_n(x) = (x^n/n!)(1 + O(n^2/x))$ . But there are many applications where we need a good estimate of  $F_n(x)$  when  $n^2/x \rightarrow \infty$  while  $n/x \rightarrow 0$ ; for example,  $x$  might be  $n \log n$ . In such cases  $\tilde{F}_n(x)$  is close to  $F_n(x)$  but  $x^n/n!$  is not.

We can express  $s/y$  as a power series in  $y$  by inverting the power series expression  $sf'(s) = y$ :

$$s/y = 1 - f_2 y + (4f_2^2 - f_3)y^2/2 + (15f_2 f_3 - 30f_2^3 - f_4)y^3/6 + \dots$$

From this we can get a formal series for  $\tilde{F}_n(x)$ ,

$$\begin{aligned} \tilde{F}_n(x) &= \frac{x^n}{n!} \frac{\exp(n(s/y)(1 + f_2 s/2! + f_3 s^2/3! + \dots) - n)}{(s/y)^n} \\ &= \frac{x^n}{n!} \left( 1 + \frac{n f_2}{2} y + \frac{3n^2 f_2^2 - 12n f_2^2 + 4n f_3}{24} y^2 + O(n^3 y^3) \right). \end{aligned}$$

We can also use the formula

$$f_{n(n-k)} = \sum \frac{n^{k+k_2+k_3+\dots}}{2!^{k_2} k_2! 3!^{k_3} k_3! \dots} f_2^{k_2} f_3^{k_3} \dots,$$

where the sum is over all nonnegative  $k_2, k_3, \dots$  with  $k_2 + 2k_3 + \dots = k$ , to write

$$F_n(x) = \frac{x^n}{n!} \left( 1 + \frac{nf_2 - f_2 + O(x^{-1})}{2} y + \frac{3n^2 f_2^2 - 18nf_2^2 + 4nf_3 + 33f_2^2 - 12f_3 + O(x^{-1})}{24} y^2 + O(n^3 y^3) \right).$$

These series are not useful asymptotically unless  $ny = n^2/x$  is small. But the approximation  $\tilde{F}_n(x)$  itself is excellent, because amazing cancellations occur when we compute the ratio:

$$\frac{F_n(x)}{\tilde{F}_n(x)} = 1 - \frac{f_2}{2} y + \frac{11f_2^2 - 4f_3}{8} y^2 + O(y^3) + O(x^{-1}).$$

**Theorem.** When  $F(z) = \exp(z + f_2 z^2/2! + f_3 z^3/3! + \dots)$  and the functions  $F_n(x)$  and  $\tilde{F}_n(x)$  are defined as above, the ratio  $F_n(x)/\tilde{F}_n(x)$  can be written as a formal power series  $\sum_{i,j \geq 0} c_{ij} y^i x^{-j}$ , where  $y = n/x$  and the coefficients  $c_{ij}$  are polynomials in  $f_2, f_3, \dots$ .

The derivation just given shows that we can write  $F_n(x)/\tilde{F}_n(x)$  as a formal power series of the form  $\sum_{i,j \geq 0} a_{ij} n^i x^{-j}$ , where  $a_{ij} = 0$  when  $i > 2j$ ; the surprising thing is that we also have  $a_{ij} = 0$  whenever  $i > j$ . Therefore we can let  $c_{ij} = a_{i(i+j)}$ .

To prove the theorem, we let  $R(z) = 1 + R_1 z + R_2 z^2 + \dots$  stand for the terms neglected in our approximation:

$$F(z)^x = e^{xf(s) - xsf'(s)} \left( 1 + \frac{n}{s} \frac{z}{1!} + \frac{n^2}{s^2} \frac{z^2}{2!} + \frac{n^3}{s^3} \frac{z^3}{3!} + \dots \right) R(z).$$

The coefficient of  $z^n$  is

$$F_n(x) = \tilde{F}_n(x) \left( 1 + R_1 s + \frac{n-1}{n} R_2 s^2 + \frac{(n-1)(n-2)}{n^2} R_3 s^3 + \dots \right);$$

so the ratio  $F_n(x)/\tilde{F}_n(x)$  is equal to

$$\sum_{k \geq 0} \frac{n^k}{n^k} R_k s^k = \sum_{j, k \geq 0} (-n)^{-j} \begin{bmatrix} k \\ k-j \end{bmatrix} R_k s^k = \sum_j (-n)^{-j} P_j,$$

where  $P_j = \sum_k \begin{bmatrix} k \\ k-j \end{bmatrix} R_k s^k$  is a certain power series in  $s$  and  $x$ . The coefficients  $R_k$  are themselves power series in  $s$  and  $x$ , because we have

$$R(z) = \exp \left( x(z-s)^2 \frac{f''(s)}{2!} + x(z-s)^3 \frac{f'''(s)}{3!} + \dots \right).$$

We know from the discussion above that

$$\begin{bmatrix} k \\ k-j \end{bmatrix} = k(k-1) \dots (k-j) \sigma_j(k)$$

is a polynomial in  $k$ . Therefore we can write

$$P_j = \left[ \begin{array}{c} \vartheta \\ \vartheta - j \end{array} \right] R(z) \Big|_{z=s},$$

where  $\vartheta$  is the operator that takes  $z^k \mapsto k z^k$  for all  $k$ ; i.e.,  $\vartheta G(z) = z G'(z)$  for all power series  $G(z)$ . The theorem will be proved if we can show that  $P_j/n^j$  is a formal power series in  $y$  and  $x^{-1}$ , and if the sum of these formal power series over all  $j$  is also such a series.

Consider, for example, the simplest case  $P_0 = R(s)$ ; obviously  $P_0 = 1$ . The next simplest case is  $P_1 = \left[ \begin{array}{c} \vartheta \\ \vartheta - 1 \end{array} \right] R(z) \Big|_{z=s} = \frac{1}{2} \vartheta(\vartheta - 1) R(z) \Big|_{z=s}$ . It is easy to see that

$$\vartheta^j = z^j D^j,$$

where  $D$  is the differentiation operator  $D G(z) = G'(z)$ , because  $z^j D^j$  takes  $z^k$  into  $k^j z^k$ . Therefore

$$P_1 = \frac{1}{2} s^2 R''(s) = \frac{1}{2} x s^2 f''(s).$$

It follows that  $P_1/n = \frac{1}{2}(s/y)s f''(s)$  is a power series in  $y$ ; it begins  $\frac{1}{2}f_2 y + \frac{1}{2}(f_3 - f_2^2)y^2 + \dots$ .

Now let's consider  $P_j$  in general. We will use the fact that the Stirling numbers  $\left[ \begin{array}{c} k \\ k-j \end{array} \right]$  can be represented in the form

$$\left[ \begin{array}{c} k \\ k-j \end{array} \right] = p_{j1} \binom{k}{j+1} + p_{j2} \binom{k}{j+2} + \dots + p_{jj} \binom{k}{2j},$$

where the coefficients  $p_{ji}$  are the positive integers in the following triangular array:

$$\begin{array}{ccccccc} 1 & & & & & & \\ 2 & & 3 & & & & \\ 6 & & 20 & & 15 & & \\ 24 & & 130 & & 210 & & 105 \\ 120 & & 924 & & 2380 & & 2520 & & 945 \end{array}.$$

(This array is clearly not a convolution matrix; but the theory developed above implies that the numbers  $j! p_{ji}/(i+j)!$ , namely

$$\begin{array}{ccccccc} 1/2 & & & & & & \\ 2/3 & & 1/4 & & & & \\ 3/2 & & 1 & & 1/8 & & \\ 24/5 & & 13/3 & & 1 & & 1/16 \\ 20 & & 22 & & 85/12 & & 5/6 & & 1/32 \end{array},$$

do form the convolution matrix for the powers of  $\exp(z/2 + z^2/3 + z^3/4 + \dots)$ . The expression  $\left[ \begin{array}{c} k \\ k-j \end{array} \right] = \sum_{i=1}^j p_{ji} \binom{k}{j+i}$  was independently discovered by [Appell 1880], [Jordan 1933], and [Ward 1934]. The number of permutations of  $i+j$  elements having no fixed points and exactly  $i$  cycles is  $p_{ji}$ , an “associated Stirling number of the first kind” [Riordan 1958, section 4.4] [Comtet 1970, exercise 6.7].) It follows that

$$P_j = p_{j1} s^{j+1} \frac{R^{(j+1)}(s)}{(j+1)!} + p_{j2} s^{j+2} \frac{R^{(j+2)}(s)}{(j+2)!} + \dots + p_{jj} s^{2j} \frac{R^{(2j)}(s)}{(2j)!}.$$

Now  $R(z)$  is a sum of terms having the form

$$a_{il}x^i(z-s)^l,$$

where  $l \geq 2i$  and where  $a_{il}$  is a power series in  $s$ . Such a term contributes  $a_{il}x^i s^l p_{j(l-j)}$  to  $P_j$ ; so it contributes  $a_{il}(s/y)^j s^{l-j} x^{i-j} p_{j(l-j)}$  to  $P_j/n^j$ . This contribution is nonzero only if  $j < l \leq 2j$ . Since  $l \geq 2i$ , we have  $i \leq j$ ; so  $P_j/n^j$  is a power series in  $y$  and  $x^{-1}$ .

For a fixed value of  $j-i$ , the smallest power of  $y$  that can occur in  $P_j/n^j$  is  $y^{2i-j} = y^{j-2(j-i)}$ . Therefore only a finite number of terms of  $\sum_j P_j/(-n)^j$  contribute to any given power of  $y$  and  $x^{-1}$ . This completes the proof.

A careful analysis of the proof, and a bit of *Mathematica* hacking, yields the more precise result

$$\frac{F_n(x)}{\tilde{F}_n(x)} = \frac{1}{(1+s^2y^{-1}d_2)^{1/2}} + \frac{(s/y)^3 A}{x(1+s^2y^{-1}d_2)^{7/2}} + O(x^{-2}),$$

where  $A = \frac{1}{12}s^3y^{-1}d_2^3 - \frac{3}{4}sd_2^2 - \frac{1}{2}s^2d_2d_3 - \frac{5}{24}s^3d_3^2 + \frac{1}{3}yd_3 + \frac{1}{8}s^3d_2d_4 + \frac{1}{8}syd_4$  and  $d_k = f^{(k)}(s)$ .

**Acknowledgment.** I wish to thank Ira Gessel and Svante Janson for stimulating my interest in this subject and for their helpful comments on the first draft. Ira Gessel and Richard Brent also introduced me to several relevant references.

## References

- Appell, P. 1880. “Développement en série entière de  $(1+ax)^{1/x}$ .” *Archiv der Mathematik und Physik* **65**: 171–175.
- Bell, E. T. 1934. “Exponential numbers.” *American Mathematical Monthly* **41**: 411–419.
- Carlitz, L. 1978. “Generalized Stirling and related numbers.” *Rivista di Matematica della Università di Parma*, serie 4, **4**: 79–99.
- Cayley, A. 1859. “On the analytical forms called trees. Second part.” *Philosophical Magazine* **18**: 371–378. Reprinted in Cayley’s *Collected Mathematical Papers* **4**: 112–115.
- Comtet, Louis. 1970. *Analyse Combinatoire*. Presses Universitaires de France. (English translation, *Advanced Combinatorics*, D. Reidel, Dordrecht, 1974.)
- Garsia, Adriano M. 1973. “An exposé of the Mullin-Rota theory of polynomials of binomial type.” *Linear and Multilinear Algebra* **1**: 47–65.
- Gomes Teixeira, F. 1896. “Sur le développement de  $x^k$  en série ordonnée suivant les puissances du sinus de la variable.” *Nouvelles Annales de Mathématiques*, série 3, **15**: 270–274.
- Good, I. J. 1957. “Saddle-point methods for the multinomial distribution.” *Annals of Mathematical Statistics* **28**: 861–881.
- Gould, H. W., and Kaucký, I. 1966. “Evaluation of a class of binomial coefficient summations.” *Journal of Combinatorial Theory* **1**: 233–248.
- Graham, Ronald L., Knuth, Donald E., and Patashnik, Oren. 1989. *Concrete Mathematics*. Addison-Wesley, Reading, Massachusetts.

- Harris, Bernard, and Schoenfeld, Lowell. 1967. "The number of idempotent elements in symmetric semigroups." *Journal of Combinatorial Theory* **3**: 122–135.
- Jabotinsky, Eri. 1947. "Sur la représentation de la composition de fonctions par un produit de matrices. Application à l'itération de  $e^x$  et de  $e^x - 1$ ." *Comptes Rendus Hebdomadaires des Sciences de L'Académie des Sciences*, **224**: 323–324.
- Jabotinsky, Eri. 1953. "Representation of functions by matrices. Application to Faber polynomials." *Proceedings of the American Mathematical Society* **4**: 546–553.
- Jabotinsky, Eri. 1963. "Analytic iteration." *Transactions of the American Mathematical Society* **108**: 457–477.
- Jordan, Charles. 1933. "On Stirling's numbers." *Tôhoku Mathematical Journal* **37**: 254–278.
- Knuth, Donald E. 1992. "Two notes on notation." *American Mathematical Monthly* **99**: 403–422.
- Knuth, Donald E. and Pittel, Boris. 1989. "A recurrence related to trees." *Proceedings of the American Mathematical Society* **105**: 335–349.
- Kuczma, Marek. 1968. *Functional Equations in a Single Variable*. Polish Scientific Publishers, Warsaw.
- Lah, I. 1955. "Eine neue Art von Zahlen, ihre Eigenschaften und Anwendung in der mathematischen Statistik," *Mitteilungsblatt für Mathematische Statistik* **7**: 203–212.
- Motzkin, T. S. 1971. "Sorting numbers for cylinders and other classification numbers." *Proceedings of Symposia in Pure Mathematics* **19**: 167–176.
- Mullin, Ronald, and Rota, Gian-Carlo. 1970. "On the foundations of combinatorial theory. III. Theory of binomial enumeration." In *Graph Theory and Its Applications*, edited by Bernard Harris (Academic Press, 1970), 167–213.
- Riordan, John. 1958. *An Introduction to Combinatorial Analysis*. John Wiley & Sons, New York.
- Riordan, John. 1968. *Combinatorial Identities*. John Wiley & Sons, New York.
- Rogers, D. G. 1978. "Pascal triangles, Catalan numbers and renewal arrays." *Discrete Mathematics* **22**: 301–310.
- Roman, Steven. 1978. *The Umbral Calculus*. Academic Press, Orlando.
- Roman, Steven M., and Rota, Gian-Carlo. 1978. "The umbral calculus." *Advances in Mathematics* **27**: 95–188.
- Rota, Gian-Carlo, Kahaner, D., and Odlyzko, A. 1973. "On the foundations of combinatorial theory. VIII. Finite operator calculus." *Journal of Mathematical Analysis and Applications* **42**: 884–760. Reprinted in Rota, Gian-Carlo, *Finite Operator Calculus* (Academic Press, 1975), 7–82.
- Steffensen, J. F. 1941. "The poweroid, an extension of the mathematical notion of power." *Acta Mathematica* **73**: 333–366.
- Ward, Morgan. 1934. "The representation of Stirling's numbers and Stirling's polynomials as sums of factorials." *American Journal of Mathematics* **56**: 87–95.